

**UNITED STATES  
SECURITIES AND EXCHANGE COMMISSION**  
Washington, D.C. 20549

**FORM 6-K**

**REPORT OF FOREIGN PRIVATE ISSUER PURSUANT TO RULE 13a-16 OR 15d-16  
UNDER THE SECURITIES EXCHANGE ACT OF 1934**

**FOR THE MONTH OF NOVEMBER 2025**

**COMMISSION FILE NUMBER 001-39081**

**BioNTech SE**

(Translation of registrant's name into English)

**An der Goldgrube 12  
D-55131 Mainz  
Germany  
+49 6131-9084-0**

(Address of principal executive offices)

Indicate by check mark whether the registrant files or will file annual reports under cover Form 20-F or Form 40-F: Form 20-F  Form 40-F

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(1):

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(7):

**DOCUMENTS INCLUDED AS PART OF THIS FORM 6-K**

On November 11, 2025, BioNTech SE (the “Company”) hosted an edition of the Company’s Innovation Series R&D Day, providing an overview of the Company’s strategy and clinical progress across its pipeline. The presentation is attached hereto as Exhibit 99.1.

**SIGNATURE**

Pursuant to the requirements of the Exchange Act, the registrant has duly caused this report to be signed on its behalf by the undersigned, thereunto duly authorized.

**BioNTech SE**

By: /s/ Ramon Zapata-Gomez  
Name: Ramon Zapata-Gomez  
Title: Chief Financial Officer

By: /s/ Dr. Sierk Poetting  
Name: Dr. Sierk Poetting  
Title: Chief Operating Officer

Date: November 12, 2025

**EXHIBIT INDEX**

| <u>Exhibit</u> | <u>Description of Exhibit</u> |
|----------------|-------------------------------|
| 99.1           | <a href="#">Presentation</a>  |

# Innovation Series: R&D Day 2025

November 11<sup>th</sup>, 2025

BIONTECH



# 1 Introductory Remarks

Douglas Maffei, PhD,  
Vice President, Strategy and  
Investor Relations

BIONTECH

## This Slide Presentation Includes Forward-Looking Statements

This presentation contains forward-looking statements within the meaning of the Private Securities Litigation Reform Act of 1995, as amended, including, but not limited to, statements concerning: BioNTech's expected revenues and net profit/(loss) related to sales of BioNTech's COVID-19 vaccine, referred to as COMIRNATY where approved for use under full or conditional marketing authorization, in territories controlled by BioNTech's collaboration partners, particularly for those figures that are derived from preliminary estimates provided by BioNTech's partners; the rate and degree of market acceptance of BioNTech's COVID-19 vaccine and, if approved, BioNTech's investigational medicines; expectations regarding anticipated changes in COVID-19 vaccine demand, including changes to the ordering environment and expected regulatory recommendations to adapt vaccines to address new variants or sublineages; the initiation, timing, progress, results, and cost of BioNTech's research and development programs, including BioNTech's current and future preclinical studies and clinical trials, including statements regarding the expected timing of initiation, enrollment, and completion of studies or trials and related preparatory work and the availability of results, and the timing and outcome of applications for regulatory approvals and marketing authorizations; BioNTech's expectations regarding potential future commercialization in oncology, including goals regarding timing and indications; the targeted timing and number of additional potentially registrational trials, and the registrational potential of any trial BioNTech may initiate; discussions with regulatory agencies; BioNTech's expectations with respect to intellectual property; the impact of BioNTech's collaboration and licensing agreements, including BioNTech's partnership with BMS; BioNTech's planned acquisition of CureVac; the development, nature and feasibility of sustainable vaccine production and supply solutions; the deployment of AI across BioNTech's preclinical and clinical operations; BioNTech's expectations with respect to tariff policy; BioNTech's estimates of revenues, research and development expenses, selling, general and administrative expenses, and capital expenditures for operating activities; BioNTech's expectations regarding upcoming payments relating to litigation settlements; BioNTech's expectations for upcoming scientific and investor presentations; and BioNTech's expectations of net profit / (loss). In some cases, forward-looking statements can be identified by terminology such as "will," "may," "should," "expects," "intends," "plans," "aims," "anticipates," "believes," "estimates," "predicts," "potential," "continue," or the negative of these terms or other comparable terminology, although not all forward-looking statements contain these words.

The forward-looking statements in this presentation are based on BioNTech's current expectations and beliefs of future events and are neither promises nor guarantees. You should not place undue reliance on these forward-looking statements because they involve known and unknown risks, uncertainties, and other factors, many of which are beyond BioNTech's control, and which could cause actual results to differ materially and adversely from those expressed or implied by these forward-looking statements. These risks and uncertainties include, but are not limited to: the uncertainties inherent in research and development, including the ability to meet anticipated clinical endpoints, commencement and/or completion dates for clinical trials, projected data release timelines, regulatory submission dates, regulatory approval dates and/or launch dates, as well as risks associated with preclinical and clinical data, including the data discussed in this release, and including the possibility of unfavorable new preclinical, clinical or safety data and further analyses of existing preclinical, clinical or safety data; the nature of the clinical data, which is subject to ongoing peer review, regulatory review and market interpretation; BioNTech's pricing and coverage negotiations regarding its COVID-19 vaccine with governmental authorities, private health insurers and other third-party payors; the future commercial demand and medical need for initial or booster doses of a COVID-19 vaccine; the impact of tariffs and escalations in trade policy; competition from other COVID-19 vaccines or related to BioNTech's other product candidates, including those with different mechanisms of action and different manufacturing and distribution constraints, on the basis of, among other things, efficacy, cost, convenience of storage and distribution, breadth of approved use, side-effect profile and durability of immune response; the timing of and BioNTech's ability to obtain and maintain regulatory approval for its product candidates; the ability of BioNTech's COVID-19 vaccines to prevent COVID-19 caused by emerging virus variants; BioNTech's and its counterparties' ability to manage and source necessary energy resources; BioNTech's ability to identify research opportunities and discover and develop investigational medicines; the ability and willingness of BioNTech's third-party collaborators to continue research and development activities relating to BioNTech's development candidates and investigational medicines; the impact of COVID-19 on BioNTech's development programs, supply chain, collaborators and financial performance; unforeseen safety issues and potential claims that are alleged to arise from the use of products and product candidates developed or manufactured by BioNTech; BioNTech's and its collaborators' ability to commercialize and market BioNTech's COVID-19 vaccine and, if approved, its product candidates; BioNTech's ability to manage its development and related expenses; regulatory and political developments in the United States and other countries; BioNTech's ability to effectively scale its production capabilities and manufacture its products and product candidates; risks relating to the global financial system and markets; and other factors not known to BioNTech at this time. You should review the risks and uncertainties described under the heading "Risk Factors" in BioNTech's Report on Form 6-K for the period ended September 30, 2025, and in subsequent filings made by BioNTech with the SEC, which are available on the SEC's website at [www.sec.gov](http://www.sec.gov). These forward-looking statements speak only as of the date hereof. Except as required by law, BioNTech disclaims any intention or responsibility for updating or revising any forward-looking statements contained in this presentation in the event of new information, future developments or otherwise.

Furthermore, certain statements contained in this presentation relate to or are based on studies, publications, surveys and other data obtained from third-party sources and BioNTech's own internal estimates and research. While BioNTech believes these third-party sources to be reliable as of the date of this presentation, it has not independently verified, and makes no representation as to the adequacy, fairness, accuracy or completeness of, any information obtained from third-party sources. In addition, any market data included in this presentation involves assumptions and limitations, and there can be no guarantee as to the accuracy or reliability of such assumptions. While BioNTech believes its own internal research is reliable, such research has not been verified by any independent source. In addition, BioNTech is the owner of various trademarks, trade names and service marks that may appear in this presentation. Certain other trademarks, trade names and service marks appearing in this presentation are the property of third parties. Solely for convenience, the trademarks and trade names in this presentation may be referred to without the ® and TM symbols, but such references should not be construed as any indicator that their respective owners will not assert, to the fullest extent under applicable law, their rights thereto.

**An abbreviation directory of defined terms can be found at the end of the presentation.**

## Innovation Series R&D Day 2025 Agenda

|   |   |   |
|---|---|---|
| 1 | <b>Introductory Remarks</b>   | Douglas Maffei, PhD,<br>Vice President, Strategy and Investor Relations   |
| 2 | <b>BioNTech's Unique Approach to Innovation</b>   | Prof. Uğur Şahin<br>Co-founder and Chief Executive Officer  |
| 3 | <b>BioNTech's Differentiated Clinical Strategy to Advance the Treatment of Solid Tumors</b> | Prof. Özlem Türeci, M.D.<br>Co-founder and Chief Medical Officer  |
| 4 | <b>Establishing Punitamig<sup>1</sup> in Foundational Tumor Types</b>                       | Prof. Ilhan Celik, M.D.<br>Vice President, Clinical Development<br>Michael Wenger, M.D.<br>Vice President, Clinical Development |
| 5 | <b>Innovating Early-Stage Cancer Treatment with mRNA Cancer Immunotherapies</b>             | Prof. Özlem Türeci, M.D.<br>Co-founder and Chief Medical Officer  |
| 6 | <b>BioNTech's Path to Value Creation</b>  | Ramón Zapata<br>Chief Financial Officer   |
| 7 | <b>Q&amp;A Panel Discussion</b>   | All Speakers<br>Annemarie Hanekamp<br>Chief Commercial Officer  |

<sup>1</sup> Partnered with Bristol Myers Squibb.



2

## BioNTech's Unique Approach to Innovation

Prof. Uğur Şahin  
CEO and Co-founder

BIONTECH











A microscopic view of various cells, including some with prominent nuclei and others that are more spherical and textured. The background is a dark, teal color with a subtle gradient.

BIONTECH


**Translating Science into Survival**  
Building a Global Immunotherapy Powerhouse

---


BioNTech – Disruptive Tech-Bio Company with Pioneering Technologies  
Developed Through Full AI Integration

|  |  |  |  |                  |  |
|--|--|--|--|------------------|--|
| <b>Multiplatform oncology company</b>  |  | <b>Infectious diseases pipeline</b>  |  |                  |  |
| <b>16</b> Clinical programs  | <b>&gt;20</b> Ongoing Phase 2 or 3 trials  | <b>7</b> Clinical programs in high unmet need indications                                    |  |                  |  |
|  Bristol Myers Squibb |  Genmab                           |  DualityBio |  MedLink Therapeutics | Gates Foundation |  CEPI |
|  OncoC4               |  Genentech                        |  REGENERON  |  |                  |  |
| <b>COVID-19 vaccine global impact</b>  | <b>Leader in integrated AI capabilities</b>  | <b>In-house manufacturing</b>  |  |                  |  |
| <b>5</b> Billion doses distributed   |  InstaDeep™<br>a BioNTech Company | <b>4</b> Platforms including individualized mRNA and bispecific antibodies                   |  |                  |  |
|  Pfizer               |  |  |  |                  |  |

— Root Cause of Cancer Treatment Failure




— We Are Uniquely Positioned to Combine Approaches to Transform Cancer Care








1. Synergistic potential.

— We Are Uniquely Positioned to Combine Approaches to Transform Cancer Care



1. Synergistic potential.


## Prioritized Immunomodulator Pipeline

| Pumitamig <sup>1</sup>   | Gotistobart <sup>2</sup>  | BNT314/<br>GEN1059 <sup>3</sup>  |
|--|---|--|
|   |   |   |
| <p>PD-L1 expression or upregulation in tumors may enrich <b>VEGF neutralization</b> into the TME which <b>inhibits angiogenesis</b>.</p>   | <p>Monospecific antibody with <b>optimized Fc</b> targeting <b>CTLA-4</b> and <b>selectively depleting tumor-infiltrating Tregs</b> in the TME but not in the periphery due to a pH driven mechanism.</p> | <br> |
| <p><b>Clinical status</b></p> <ul style="list-style-type: none"> <li>• <b>Registrational trials</b> ongoing in 1L SCLC, NSCLC, TNBC and initiating in CRC, gastric</li> <li>• 12+ studies combining with chemotherapy</li> <li>• 10+ novel combinations</li> </ul> | <p><b>Clinical status</b></p> <ul style="list-style-type: none"> <li>• <b>Ph3</b> in 2L+ sqNSCLC</li> <li>• Ph2 in PROC</li> <li>• Ph1/2 in mCRPC</li> <li>• Ph1/2 in multiple solid tumors</li> </ul>    | <p><b>Clinical status</b></p> <p>Phase 1, Phase 1/2, exploratory trials ongoing</p> <p>Exploratory exercise: More novel next-gen IO molecules to come</p>                  |

1. Partnered with Bristol Myers Squibb; 2. Partnered with OncoC4; 3. Partnered with Genmab.


## Pumitamig<sup>1</sup>: PD-L1 x VEGF-A Bispecific Antibody

Pumitamig is an investigational bispecific antibody, targeting both PD-L1 and VEGF-A. Binding to PD-L1 is intended to restore effector T-cell function and localize VEGF-A neutralization within the TME, reversing the negative impact of VEGF signaling on immune cell infiltration and activation and normalizing tumor vasculature, leading to tumor growth inhibition.




Miao X et al. AACR 2025. Poster #6061; 1. Partnered with Bristol Myers Squibb

## Pumitamig<sup>1</sup>: Potent VEGF-A Neutralization and PD-1/PD-L1 Checkpoint Blockade




Data on file. 1. Partnered with Bristol Myers Squibb; 2. Luciferase-based reporter assays are commercially available from Promega.

# Pumitami<sup>1</sup>: Combined Effect of Anti-Angiogenesis and Checkpoint Blockade Through One Molecule Leads to Immune Invigoration



Data on file. 1. Partnered with Bristol Myers Squibb


## Pumitamidg<sup>1</sup> is More than the Sum of Two Monospecific Antibodies



1. Partnered with Bristol Myers Squibb

# VEGF-A / Bispecific Antibody Complex Formation is a Function of Antibody-to-VEGF-A Molar Ratio

Complex formation occurs at optimal antibody-to-VEGF-A molar ratios in a bell-shaped response curve




Data on file.


# Pumitamid<sup>1</sup> / VEGF-A Complexation Leads to Enhanced Binding Affinity to PD-L1

At equimolar pumitamid-to-VEGF-A ratio, pumitamid shows increased On-rate and decreased Off-rate

A. Normalized SPR sensorgrams



B. Isoaffinity graph




C.  $K_D$  values and association ( $k_a$ ) /dissociation ( $k_{dis}$ ) rates for pumitamid

| Pumitamid:VEGF-A molar ratio | $K_D$ (Binding affinity) [M]               | $k_a$ [ $M^{-1}s^{-1}$ ]              | $k_{dis}$ [ $s^{-1}$ ]                   | Fold differences |
|------------------------------|--|---------------------------------------|--|------------------|
| Pumitamid only               | $4.915 \times 10^{-10}$                    | $1.651 \times 10^6$                   | $8.114 \times 10^{-4}$                   |                  |
| 100:1                        | $3.468 \times 10^{-10}$                    | $2.063 \times 10^6$                   | $7.153 \times 10^{-4}$                   | 1.4              |
| 10:1                         | $1.317 \times 10^{-10}$                    | $1.684 \times 10^6$                   | $2.217 \times 10^{-4}$                   | 3.7              |
| 5:1                          | $1.158 \times 10^{-10}$                    | $1.977 \times 10^6$                   | $2.289 \times 10^{-4}$                   | 4.2              |
| 2:1                          | $0.4029 \times 10^{-10}$                   | $6.299 \times 10^6$                   | $2.538 \times 10^{-4}$                   | 12.2             |
| 1:1                          | <b><math>0.1537 \times 10^{-10}</math></b> | <b><math>1.011 \times 10^7</math></b> | <b><math>1.555 \times 10^{-4}</math></b> | <b>32</b>        |


**32-fold enhanced binding affinity of pumitamid to immobilized PD-L1 by complexation of pumitamid with equimolar VEGF-A, indicative of cooperative binding**

Data on file. 1. Partnered with Bristol Myers Squibb

# Rapid anti-VEGF-A Dependent Internalisation of Pumitamig<sup>1</sup> Upon Binding to PD-L1<sup>+</sup> Cells



PD-L1<sup>+</sup>A549 cells treated with antibodies which were pre-incubated with an equimolar ratio of VEGF-A; red fluorescence indicates internalization




Rapid PD-L1 internalization may contribute to enhanced T cell activation in the presence of Pumitamig / VEGF-A complexes


Data on file. 1. Partnered with Bristol Myers Squibb

# Pumitamig<sup>1</sup>: Synergistic Targeting of PD-L1 and VEGF-A


## Pumitamig characteristics: combined tumor targeting<sup>2</sup>



### Selected NSCLC IHC<sup>3</sup>



VEGF-A



PD-L1


### Bispecific MOA

Targeting of PD-1/PD-L1 blockade to VEGF-A high tumors

Targeting of VEGF-A neutralization to PDL1 high tumors


1. Partnered with Bristol Myers Squibb; 2. Khan KA Nat Rev Clin Oncol 2018; 3. IHC data: Human Protein Atlas.

— Differentiation of Punitamig<sup>1</sup> by Binding to PD-L1 Aims Targeting to Tumor Site

|                                 | Blocking of PD-1/PD-L1 signaling | Neutralization of VEGF | Cooperative effect linking PD-L1 and VEGF binding | TME Targeting by anti-PD-L1 | <b>Punitamig<sup>1</sup></b><br><b>Dual targeting of TME</b><br>VEGF targeted PD-L1 inhibition<br>Anti-VEGF-A<br><br>Anti-PD-L1<br>PD-L1 targeted VEGF neutralization |
|---------------------------------|----------------------------------|------------------------|---|-----------------------------|--|
| Punitamig PD-L1/VEGF Bispecific | YES                              | YES                    | YES   | YES                         |  |
| PD-1/VEGF Bispecific            | YES                              | YES                    | YES   | NO                          |  |
| PD(L)1 + VEGF Monospecific      | YES                              | YES                    | NO  | NO                          |  |

1. Partnered with Bristol Myers Squibb.


Pumitamig<sup>1</sup> May Drive Clinical Benefit Irrespective of PD-L1 Status




# Pumitami<sup>1</sup> in Combination with Chemo Shows Efficacy in EGFRm NSCLC Irrespective of PD-L1 Status

## Phase 2 study\* of pumitami<sup>1</sup> + carbo/pem in EGFRmut NSCLC post EGFR TKI

Adapted from Wu YL et al. ESMO 2024 1255MO



Broad Combination Strategy Across Indications Aiming to Establish Next-Generation IO-Backbone








1. NCI SEER <https://training.seer.cancer.gov/index.html>. 2. US incidence source: NIH and American Cancer Society data EU incidence source: European Cancer Information System

Pumitamig<sup>1</sup>: Executing a Parallel Three-Wave Strategy to Build a Proprietary IO Franchise




Partnered with 1. Bristol Myers Squibb.

## Prioritized Immunomodulator Pipeline

| Pumitamig <sup>1</sup>   | Gotistobart <sup>2</sup>  | BNT314/<br>GEN1059 <sup>3</sup>   |
|--|---|---|
|  <p>Anti-VEGF A<br/>Inert Fc (LALA)<br/>Anti-PD-L1 VHH</p>  |  <p>Anti-CTLA4<br/>Optimized Fc</p>   |  <p>Anti-4-1BB Anti-EpCAM<br/>Inert Fc</p>   |
| <p>PD-L1 expression or upregulation in tumors may enrich <b>VEGF neutralization</b> into the TME which <b>inhibits angiogenesis</b>.</p>   | <p>Monospecific antibody with <b>optimized Fc</b> targeting <b>CTLA-4</b> and <b>selectively depleting tumor-infiltrating Tregs</b> in the TME but not in the periphery due to a pH driven mechanism.</p> |  <p>Anti-PD-1<br/>CD38</p>  <p>Anti-TIGIT<br/>PVRIG</p> |
| <p><b>Clinical status</b></p> <ul style="list-style-type: none"> <li>• <b>Registrational trials</b> ongoing in 1L SCLC, NSCLC, TNBC and initiating in CRC, gastric</li> <li>• 12+ studies combining with chemotherapy</li> <li>• 10+ novel combinations</li> </ul> | <p><b>Clinical status</b></p> <ul style="list-style-type: none"> <li>• <b>Ph3</b> in 2L+ sqNSCLC</li> <li>• Ph2 in PROC</li> <li>• Ph1/2 in mCRPC</li> <li>• Ph1/2 in multiple solid tumors</li> </ul>    | <p><b>Clinical status</b></p> <p>Phase1, Phase 1/2, exploratory trials ongoing</p> <p>Exploratory exercise: More novel next-gen IO molecules to come...</p>   |

1. Partnered with Bristol Myers Squibb; 2. Partnered with OncoC4; 3. Partnered with Genmab.

## Anti-CTLA4 as a Target for Cancer Immunotherapy




## Gotistobart<sup>1</sup> Differentiated Mechanism with Potential to Become Best-in-Class Anti-CTLA-4 Antibody

Avoiding lysosomal degradation of CTLA-4 for safer and more effective immunotherapy may lead to uncoupling cancer therapeutic effect from immunotherapy-related adverse effects

### Gotistobart<sup>1</sup> designed to:


- Allow regular recycling and enrichment of antibody and CTLA-4 molecule
- Selectively kill Tregs in the tumor microenvironment
- Improve therapeutic index (efficacy/toxicity ratio)
- Enhance anti-tumor immunity
- Allow prolonged, repeated dosing




Liu Y. et al. SITC 2021 # 231; Du et al. Cell Res. 2018 Apr; 28(4): 416–432; Du et al. Cell Res. 2018 Apr; 28(4): 433–447.

1. Partnered with OncoC4.

Preclinical Data Demonstrate Improved Therapeutic Index of Gotistobart<sup>1</sup>



Zhang *et al.* Cell Research 2019




Data on file.

1. Partnered with OncoC4.


# CD3+, CD4+ and FoxP3+ Treg Concentrations Decrease with Gotistobart<sup>1</sup> Treatment

PRESERVE-001 Phase 1/2 Trial: gotistobart monotherapy treatment in multiple tumor types, including sarcoma, pancreatic cancer, melanoma and ovarian cancer




1. Partnered with OncoC4. Data on file

# Pivotal Development of Gotistobart<sup>1</sup> in 2L Squamous Non-Small Cell Lung Cancer




— We are Uniquely Positioned to Combine Approaches to Transform Cancer Care



1. Synergistic potential.


## mRNA-Technology Toolbox



Holtkamp et al. Blood 2006; Kuhn et al. Gene Therapy 2010; Sahin, Türeci & Kariko Nat Drug Discovery 2014; Vogel et al. Mol Therapy 2018; Beisert et al. Mol Therapy 2020.



# Nanoparticulate mRNA Vaccines

## BNT111 mRNA LPX Vaccine




**Systemic Targeting to Lymphoid Dendritic Cells**

Kranz, Diken et al., Nature 2016



## Nanoparticulate mRNA-LPX Vaccines for Cancer Immunotherapy




### Systemic RNA delivery to dendritic cells exploits antiviral defence for cancer immunotherapy

Lena M. Kranz<sup>1,2\*</sup>, Mustafa Diken<sup>1,3\*</sup>, Heinrich Haas<sup>3</sup>, Sebastian Kreiter<sup>1,3</sup>, Carmen Logigal<sup>4</sup>, Kerstin C. Reuter<sup>1</sup>, Martin Meng<sup>1</sup>, Daniel Fritz<sup>2</sup>, Fulvia Vascotto<sup>3</sup>, Hossam Hefesh<sup>3</sup>, Christian Grunwitz<sup>2,3</sup>, Mathias Vormehr<sup>2,3</sup>, Yves Hüsemann<sup>3</sup>, Abderraouf Selmi<sup>1,2</sup>, Andreas N. Kuhn<sup>3</sup>, Janina Buck<sup>3</sup>, Evelyn Derhovanessian<sup>3</sup>, Richard Rae<sup>1</sup>, Sebastian Attig<sup>2,3</sup>, Jan Diekmann<sup>3</sup>, Robert A. Jabulowsky<sup>3</sup>, Sandra Heesch<sup>3</sup>, Jessica Hassel<sup>3</sup>, Peter Langguth<sup>4</sup>, Stephan Grabbe<sup>4</sup>, Christoph Huber<sup>1,3</sup>, Özdem Türeci<sup>5</sup> & Ugur Sahin<sup>1,2,3\*</sup>

- RNA-LPX vaccine targeting dendritic cells
- TLR7 driven adjuvant effect
- Type-I Interferon driven innate & adaptive immune stimulation
- Universally applicable for almost all type of tumor antigens
- Overcomes tolerance against self-antigens
- Excellent immunogenicity *in vivo*
- Preclinical activity against advanced tumors

— We are Uniquely Positioned to Combine Approaches to Transform Cancer Care



1. Synergistic potential.

## ADC Innovation Cycle is Just Beginning

### BioNTech is driving the development of next-generation ADCs

#### Distinguished ADC linker technology

- Stability improving safety profile
- Higher efficacy


#### Novel mechanisms of actions

- Tumor specific activation
- Improved and novel payloads

#### Novel targets and novel epitopes


- Targeting broader spectrum of tumors
- Higher specificity

#### BioNTech is developing ADCs against novel targets



Our deep understanding of ADC targets and immunology distinctively positions us to consolidate and maximize the substantial therapeutic window offered by the next-gen ADC technology


— ADC Driven Mechanisms May Synergize with Other Immune Mechanisms



**Targeted Tumor Cell Killing**  
**Bystander Tumor Cell Killing**  
**Immunogenic Cell Death**  
**Promoting Innate and Adaptive Immunity**  
**Opportunity for Synergy with Immunotherapy Compounds**


Adapted from Fucà G, et al.  
Int J Gynecol Cancer. 2024 Nov

— We are Uniquely Positioned to Combine Approaches to Transform Cancer Care





1. Synergistic potential.

# Pumitami<sup>1</sup> in Combination with ADCs Showed Superior Anti-Tumor Activity Compared with Each Treatment Alone in Pre-Clinical Tumor Models




Pumitami refers to the total mouse surrogate Pumitami antibody that binds murine VEGF-A/PlD-L1 targets; Data shown represents mean±SEM.  
 Statistical significance testing was performed comparing treatment groups with PBS/saline (\*, \*\*, \*\*\*, \*\*\*\*), with ADC-monotherapy (\*, \*\*, \*\*\*, \*\*\*\*) or with (m)Pumitami-monotherapy (\*, \*\*, \*\*\*, \*\*\*\*).  
 Source: <https://doi.org/10.1158/1538-7445.AM2025-648>

## Pumitamig<sup>1</sup> in Combination with BNT3212 Showed Synergistic Anti-Tumor Activity



<sup>1</sup>. Partnered with Bristol Myers Squibb. A375 tumor cells and human PBMCs were co-implanted subcutaneously in B-NDG B2M KO Plus mice.

## BNT3213 and Punitamig<sup>1</sup> Showed Superior Anti-Tumor Activity Compared with Each Treatment Alone in Pre-Clinical Tumor Models




<sup>1</sup> Partnered with Bristol Myers Squibb; A375 tumor cells and human PBMCs were co-implanted subcutaneously in NOD-SCID mice.

# Combined Checkpoint Inhibition and 4-1BB Agonism of Pumitamig<sup>1</sup> plus BNT314/GEN1059<sup>2</sup> In Vitro

Pumitamig<sup>1</sup> + BNT314/GEN1059<sup>2</sup> enhanced expression of cytotoxic markers and cytotoxic activity of CD8<sup>+</sup> T cells compared to single-agent treatments *in vitro*

Imle et al. SITC 2025, P652



1. Partnered with Bristol Myers Squibb; 2. Partnered with Genmab. \*\*\*, P<0.001; \*\*, P<0.01; \*, P<0.05

## Novel Combination Trials Across Multiple Tumor Types

| Combination Partners |                          | Indications                   |   |
|----------------------|--------------------------|-------------------------------|---|
| Next Gen IO + ADC    | Pumitamig <sup>1</sup>   | + T-Pam <sup>2</sup>          | HR+ HER2-low, ultra-low/null BC or TNBC                       |
|                      |                          | + BNT324/DB-1311 <sup>2</sup> | NSCLC, SCLC, HCC, melanoma, HNSCC, PROC                       |
|                      |                          | + BNT325/DB-1305 <sup>2</sup> | TNBC, NSCLC, OC   |
|                      |                          | + BNT326/YL202 <sup>3</sup>   | NSCLC, EGFRm NSCLC, HER2-neg BC, melanoma, other solid tumors |
|                      |                          | + BNT3212                     | Multiple solid tumors   |
| Next Gen IO + IO     | Pumitamig <sup>1</sup>   | + BNT314/GEN1059 <sup>4</sup> | MSS-CRC   |
|                      |                          | + BNT3213                     | HCC <sup>6</sup>  |
| Next Gen IO + mRNA   | Pumitamig <sup>1</sup>   | + BNT116                      | NSCLC   |
|                      | Gotistobart <sup>5</sup> | + BNT116                      | NSCLC   |
| mRNA + ADC           | BNT116                   | + BNT324/DB-1311 <sup>2</sup> | NSCLC   |
|                      |                          | + BNT326/YL202 <sup>3</sup>   | NSCLC   |

Partnered with: 1. Bristol Myers Squibb; 2. DualityBio ; 3. MedLink; 4. Genmab; 5. Onco C4; 6. Trial ongoing in China.

# BioNTech: Advancing Tomorrow's Personalized Precision Medicine with Integrated Capabilities Under One Roof

## Fully-integrated tech-bio company



Deep genomics & immunology expertise to analyze patient data



Individualized treatment platforms to address inter-individual variability




AI-infused & digitally-integrated target & drug discovery and development



Automated in-house manufacturing to serve patients on time and globally

## Capabilities to build tomorrow's personalized precision medicines



**BioNTech Operating  
from Position of Strength**

**2026**

Key Areas of Focus

**1**

**Combination Therapy Momentum**

Anticipate additional datasets from novel-novel combination trials with pumitamidg

**2**

**Modalities to Disease Areas**

2026 marks BioNTech's movement to a focused disease area specific approach

**3**

**Late-Stage Acceleration**

Expect key late-stage data readouts for initial wave of oncology assets

BIONTECH




3

## BioNTech's Differentiated Clinical Strategy to Advance the Treatment of Solid Tumors

Prof. Özlem Türeci, M.D.  
Chief Medical Officer and Co-founder

BIONTECH

## BioNTech's Current Priority Programs









1. Synergistic potential. Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. Genentech, a member of the Roche Group; 6. Genmab; 7. OncoC4.

# Pumitamig: Executing a Parallel Three-Wave Strategy to Build a Proprietary IO Franchise

| Establish  | Expand  | Elevate  |
|--|---|--|
| <p><b>SCLC</b></p> <ul style="list-style-type: none"> <li>1L Ph3 (Global) <b>ROSETTA</b> LUNG-01</li> <li>2L Ph3 (China)</li> <li>1L/2L Ph2 (Global)</li> </ul> <p><b>NSCLC</b></p> <ul style="list-style-type: none"> <li>1L Ph2/3 (Global) <b>ROSETTA</b> LUNG-02</li> <li>2L Ph2 (Global)</li> <li>2L EGFRmut Ph2 (China)</li> <li>IIT neoadjuvant (China)</li> </ul> <p><b>TNBC</b></p> <ul style="list-style-type: none"> <li>1L Ph3 trial (Global) <b>ROSETTA</b> TNBC-01</li> <li>1L Ph3 (China)</li> </ul> | <p><b>Registrational-Intent</b></p> <ul style="list-style-type: none"> <li>1L Gastric Ph2/3 (Global) <b>ROSETTA</b> GASTRIC-004</li> <li>1L CRC Ph2/3 (Global) <b>ROSETTA</b> CRC-002</li> </ul> <p><b>Signal-Seeking</b></p> <ul style="list-style-type: none"> <li>1L PDAC Ph2 (Global) <b>ROSETTA</b> PDAC-001</li> <li>1L PDAC Ph2 (China)</li> <li>1L GBM Ph2 (Global) <b>ROSETTA</b> GBM-100</li> <li>1L GBM Ph2 (China)</li> <li>1L CRC Ph2 (China)</li> <li>1L HCC Ph2 (China)</li> <li>1L MPM Ph2 (China)</li> <li>1L NEN Ph2 (China)</li> <li>HNSCC, RCC, CC, PROC, EC, Melanoma Ph1/2 (China)</li> </ul> | <ul style="list-style-type: none"> <li>Combining with our ADCs targeting <ul style="list-style-type: none"> <li>HER2</li> <li>TROP2</li> <li>B7H3</li> <li>HER3</li> <li>EGFR x HER3</li> <li>Novel targets</li> </ul> </li> <li>Exploring potential synergies with our IO agents <ul style="list-style-type: none"> <li>EpCam x 4-1BB</li> <li>TIGIT x PVRIG</li> <li>mRNA cancer immunotherapy</li> </ul> </li> </ul> <p><b>Potential New Standards of Care</b><br/>10+ Novel-Novel Combinations</p> |
| <p><b>Broad Pan-Tumor Applicability With Standard-of-Care Chemotherapy</b><br/>12+ Studies Exploring Pumitamig<sup>1</sup> in 10+ New Indications</p>  |   |  |
| <p><b>Foundational Registrations</b><br/>Registrational Trials with Pumitamig<sup>1</sup> Ongoing in 3 High-Impact Tumors</p>  |   |  |

Partnered with 1. Bristol Myers Squibb.

# Clinical Stage ADC Program

|   | T-Pam <sup>1</sup>   | BNT324/DB-1311 <sup>1</sup>   | BNT325/DB-1305 <sup>1</sup>   | BNT326/YL202 <sup>2</sup>   | BNT329  | BNT3212  |            |                |                |             |              |                 |  |
|---|--|---|---|---|---|--|------------|----------------|----------------|-------------|--------------|-----------------|--|
|   |   |    |    |   |  |   |            |                |                |             |              |                 |  |
|   | Target: HER2<br>Payload: topo I inhibitor  | Target: B7H3<br>Payload: topo I inhibitor   | Target: TROP2<br>Payload: topo I inhibitor  | Target: HER3<br>Payload: topo I inhibitor   | Target: CA19-9<br>Payload: topo I inhibitor   | Target: EGFRxHER3<br>Payload: topo I inhibitor                                       |            |                |                |             |              |                 |  |
|   | <b>Clinical status</b><br>• 1,100+ patients dosed<br>• Ph3: HR+HER2-low mBC<br>• Ph1/2: multiple solid tumors (EC cohort fully recruited)<br>• Ph1/2: pumitamig combo (HR+ and HR- HER2+, low and null BC) | <b>Clinical status</b><br>• 600+ patients dosed<br>• Ph1/2: multiple solid tumors<br>• Ph1/2: combo with pumitamig and BNT325 | <b>Clinical status</b><br>• 500+ patients dosed<br>• Ph1/2: multiple solid tumors<br>• Ph2: combo with pumitamig and BNT324 | <b>Clinical status</b><br>• 600+ patients dosed<br>• Ph1/2: multiple solid tumors<br>• Ph1/2: combo with pumitamig (NSCLC, other solid tumors). | <b>Clinical status</b><br>• Ph1/2: multiple solid tumors                            | <b>Clinical status</b><br>• Ph1/2: multiple solid tumors<br>• Ph1/2: pumitamig combo |            |                |                |             |              |                 |  |
|   | <i>More novel next-gen ADCs to come...</i>   |   |   |   |   |  |            |                |                |             |              |                 |  |
| <b>Expression level by indication<sup>3</sup></b> | <b>Target</b>  | <b>NSCLC</b>  | <b>SCLC</b>   | <b>HER2+ BC</b>   | <b>HR+ BC</b>   | <b>TNBC</b>  | <b>CRC</b> | <b>Gastric</b> | <b>Ovarian</b> | <b>PDAC</b> | <b>HNSCC</b> | <b>Prostate</b> | <b>Other high expression indications</b> |
| High  | HER2   |   |   |   |   |  |            |                |                |             |              |                 | Gynecologic                              |
| Medium / Low                                      | TROP2  |   |   |   |   |  |            |                |                |             |              |                 | UC, EC                                   |
| Very low / None                                   | B7H3   |   |   |   |   |  |            |                |                |             |              |                 |  |
|   | HER3   |   |   |   |   |  |            |                |                |             |              |                 | UC, BTC, EC                              |
|   | CA19-9   |   |   |   |   |  |            |                |                |             |              |                 | GBM, UC, RCC                             |
|   | EGFR   |   |   |   |   |  |            |                |                |             |              |                 |  |

**Broad ADC coverage across all relevant tumors provides optionality for selecting the most suitable therapeutic approach per indication**

1. Partnered with DualityBio; 2. Partnered with MedLink; 3. Human Protein Atlas. ADC structures shown for illustration-purpose only

— Punitamig<sup>1</sup> Extensively Studied as Monotherapy and SOC Chemo Combo

| Indication \ Regimen               | Lung cancers |              |      | Breast cancers |              | Gyn cancers |          |    | GI cancers |     |      |     | GU cancers |          | Other cancers |       |          |
|------------------------------------|--------------|--------------|------|----------------|--------------|-------------|----------|----|------------|-----|------|-----|------------|----------|---------------|-------|----------|
|                                    | NSCLC AGA-   | NSCLC EGFRin | SCLC | TNBC           | HR+/HER2- BC | Endometrial | Cervical | OC | GC/GEJ     | CRC | PDAC | HCC | RCC        | Prostate | GBM           | HNSCC | Melanoma |
| Punitamig <sup>1</sup> Monotherapy | ■            | ■            |      |                |              | ■           | ■        | ■  |            |     |      | ■   | ■          |          | ■             | ■     | ■        |
| Punitamig <sup>1</sup> + Chemo     | ■            | ■            | ■    | ■              |              |             |          |    | ■          | ■   | ■    | ■   |            |          | ■             |       |          |

■ Registrational trial   ■ Phase 1/2 trial

**Over 1,400 patients\* dosed across punitamig monotherapy and chemo combination studies**

1. Partnered with Bristol Myers Squibb \*Patient number include both China and ex-China studies, sponsored by BioNTech or partners. Punitamig is also being investigated in BTC (mono), MPM (chemo combo), and NEN (mono, chemo combo).

Single Activity of ADCs Being Explored Across Indications



Partnered with 1. DualityBio; 2. MedLink \*Patient number include both China and ex-China studies, sponsored by BioNTech or partners.


Expanding Punitamig<sup>1</sup> Opportunity with Ongoing Novel ADC Combinations

| Indication \ Regimen                                 | Lung cancers |             |      | Breast cancers |             | Gyn cancers |          |    | GI cancers |     |      |     | GU cancers |          | Other cancers |       |          |
|--|--------------|-------------|------|----------------|-------------|-------------|----------|----|------------|-----|------|-----|------------|----------|---------------|-------|----------|
|  | NSCLC AGA-   | NSCLC EGFRm | SCLC | TNBC           | HR+/HER2-BC | Endometrial | Cervical | OC | GC/GEJ     | CRC | PDAC | HCC | RCC        | Prostate | GBM           | HNSCC | Melanoma |
| Punitamig <sup>1</sup> Chemo combo                   | ■            |             | ■    | ■              |             |             |          |    | ■          | ■   | ■    |     |            |          | ■             |       |          |
| Punitamig <sup>1</sup> + T-Pam <sup>2</sup>          |              |             |      | ■              | ■           |             |          |    |            |     |      |     |            |          |               |       |          |
| Punitamig <sup>1</sup> + BNT324/DB-1311 <sup>2</sup> | ■            | ■           | ■    |                |             |             | ■        | ■  |            |     |      | ■   |            |          |               | ■     | ■        |
| Punitamig <sup>1</sup> + BNT325/DB-1305 <sup>2</sup> | ■            | ■           |      | ■              |             |             | ■        | ■  |            |     |      |     |            |          |               |       |          |
| Punitamig <sup>1</sup> + BNT326/YL202 <sup>3</sup>   | ■            | ■           |      |                | ■           |             |          |    |            |     |      |     |            |          |               |       | ■        |

■ Registrational trial   ■ Phase 1/2 trial


Partnered with 1. Bristol Myers Squibb; 2. DualityBio; 3. MediLink \*Patient number include both China and ex-China studies, sponsored by BioNTech or partners.

## Our Diverse Lung Cancer Pipeline




1. Synergistic potential; Partnered with 2 DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. OncoC4.

## BioNTech's Currently Ongoing Trials\* in Lung Cancer




Partnered with: 1. Bristol Myers Squibb; 2. OncoC4; 3. DualityBio; 4. MedLink; \*As of November 2025; \*\*being conducted in China

## Our Diverse Breast Cancer Pipeline




1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb.

# BioNTech's Currently Ongoing Trials\* in Breast Cancer




Partnered with: 1. Bristol Myers Squibb; 2. DualityBio; 3. Medilink \*As of November 2025

## Our Diverse GI Cancer Pipeline



1. Synergistic potential. Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. Genmab; 6. Genentech, a member of the Roche Group.

## BioNTech's Currently Ongoing Trials\* in GI Cancers



Partnered with: 1. Bristol Myers Squibb; 2. Genmab; 3. DualityBio; 4. MediLink.\*As of November 2025



4

## Establishing Pumitamidg in Foundational Tumor Types

Prof. Ilhan Celik, M.D.  
Vice President, Clinical Development  
Michael Wenger, M.D.  
Vice President, Clinical Development


BIONTECH

# Thoracic Cancer

BIONTECH


---

## Our Diverse Lung Cancer Pipeline



1. Synergistic potential; Partnered with 2 DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. OncoC4.

## BioNTech's Currently Ongoing Trials\* in Lung Cancer




Partnered with: 1. Bristol Myers Squibb; 2. OncoC4; 3. DualityBio; 4. MedLink; \*As of Nov 2025; \*\*being conducted in China

— Non-Small Cell Lung Cancer is One of the Highest Incidence Cancers Globally<sup>1</sup>

2030 U.S., EU4, U.K. NSCLC incidence<sup>1</sup> **~415k**

NSCLC staging distribution<sup>2</sup>



Treatment outcomes vary based on histology and PD-L1 levels in 1L NSCLC patients without actionable genomic alterations

|   | Non-squamous (~ 70%) <sup>3</sup>    | Squamous (~ 30%) <sup>3</sup>        |
|---|--------------------------------------|--------------------------------------|
| PD-L1 ≥ 50% (~ 25 - 30%) <sup>4,5</sup>   | 5-year OS: 30% (KN-189) <sup>6</sup> | 5-year OS: 23% (KN-407) <sup>7</sup> |
| PD-L1 1 - 49% (~ 30 - 40%) <sup>4,5</sup> | 5-year OS: 20% (KN-189) <sup>6</sup> | 5-year OS: 21% (KN-407) <sup>7</sup> |
| PD-L1 < 1% (~ 30 - 40%) <sup>4,5</sup>    | 5-year OS: 10% (KN-189) <sup>6</sup> | 5-year OS: 11% (KN-407) <sup>7</sup> |

1. Globocan – Cancer Tomorrow. 2. CancerMPact® 2024 Treatment Architecture EU5 and US; Note that 5-year survival reported includes all comor NSCLC population ie including with actionable genetic alterations. 3. Ganti AK, et al. JAMA Oncol. 2021 Dec; 4. Mansour MSI et al. Int J Mol Sci. 2022 Apr 19;23(9):4517; 5. Saez de Gortola, K, et al. Diagnostics 2021, 11, 1452; 6. Garassino MC, et al. J Clin Oncol. 2023 Apr 10;41(11):1992-1998; 7. Silvia Novello et al. JCO 41, 1999-2006(2023).

# Pumitamid<sup>1</sup> in Non-Small Cell Lung Cancer

## Efficacy

Efficacy observed in patients with and without driver mutations and irrespective of PD-L1 levels

## Safety Profile

Manageable safety profile with low rates of discontinuation

## Focused Execution

ROSETTA Lung-02 ongoing in both non-squamous and squamous histologies. Phase 2 part completed. Phase 3 recruiting.

| Patient Population | 1L EGFR/ALK WT PD-L1+ (TPS≥1) NSQ NSCLC<br>20 mg/kg | 2L/3L EGFR mut TKI-experienced NSQ NSCLC<br>20 mg/kg | 2L EGFR/ALK WT PD-(L)1 r/r NSQ NSCLC<br>20 mg/kg | 2L/3L EGFR-mutated TKI-experienced NSQ NSCLC<br>30 mg/kg |
|--------------------|---|--|--|--|
| N                  | 17  | 36   | 8  | 64   |
| cORR (%)           | 47.1  | 19.4   | 12.5   | 57.8   |
| DCR (%)            | 100.0   | 69.4   | 62.5   | 95.3   |
| mPFS (months)      | 13.6  | 5.5  | 6.7  | -  |
| mOS (months)       | 13.9  | 15.1   | 9.4  | -  |
| Congress           | ASCO 2024   |  | ESMO 2024  |  |

Yi-Long Wu, et. al. ESMO 2024 1255MO (pumitamid + chemotherapy)



Benchmark Data<sup>2</sup> 1L NSCLC


| Indication             | Benchmark regimen | ORR | mPFS   | mOS     | Study                    |
|------------------------|-------------------|-----|--------|---------|--------------------------|
| 1L NSCLC (PD-L1 ≥ 50%) | Pembro mono       | 46% | 7.7 mo | 26.3 mo | KEYNOTE-024 <sup>3</sup> |
| 1L NSQ NSCLC           | Pembro + chemo    | 48% | 9.0 mo | 22.0 mo | KEYNOTE-189 <sup>4</sup> |
| 1L SQ NSCLC            | Pembro + chemo    | 62% | 8.0 mo | 17.2 mo | KEYNOTE-407 <sup>5</sup> |

1. Partnered with Bristol Myers Squibb; 2. This benchmarking is not based on head-to-head trials between BioNTech's investigational candidates and other products or product candidates. Furthermore, definitive conclusions cannot be drawn from cross-trial comparisons or anticipated data, as they may be confounded by various factors, and should be interpreted with caution; 3. Reck et al, NEJM 2016; 4. Garassino et al, J Clin Oncol, 2023 5. Novello et al, J Clin Oncol, 2023

Seamless Phase 2/3 multi-site, randomized trial of punitamig in combination with chemotherapy in 1L NSCLC

**Key Inclusion Criteria**

- Treatment naïve Stage IIIB/IIIC or IV NSCLC
- RECIST 1.1 measurable disease
- ECOG PS 0 or 1
- PD-L1 all-comers



**Key Endpoints**

- Primary:** PFS (BICR), OS
- Secondary:** PFS (inv), ORR

**Benchmark Data 1L NSCLC**

| Histologies | Regimen                     | ORR | mPFS   | mOS     | Study                    |
|-------------|-----------------------------|-----|--------|---------|--------------------------|
| NSQ NSCLC   | Pembro + chemo <sup>2</sup> | 48% | 9.0 mo | 22.0 mo | KEYNOTE-189 <sup>4</sup> |
| SQ NSCLC    | Pembro + chemo <sup>3</sup> | 62% | 8.0 mo | 17.2 mo | KEYNOTE-407 <sup>5</sup> |

1. Partnered with Bristol Meyer Squibb; 2. Carboplatin + pemetrexed → pemetrexed maintenance. 3. carboplatin + paclitaxel / nab-paclitaxel. 4. Garassino et al., J Clin Oncol, 2023 5. Novello et al., J Clin Oncol, 2023; NCT06712316.

## Squamous NSCLC Remains an Area of High Unmet Need

By 2030

**55k** squamous patients start in 1L (non-AGA population)<sup>1</sup>

**~30%** continue into 2L treatment and are IO addressable

Amongst NSCLC, metastatic squamous NSCLC is seen as **#1 area of unmet need** for improving treatment amongst NSCLC<sup>2</sup>

**Limited treatment options** for patients without actionable genetic alterations in squamous NSCLC

In 2L, current chemo-based SOC shows 10 months median OS in clinical trials

<25% respond to 2L chemo-based SOC (docetaxel ± ramucirumab)

Multiple efforts failed to improve therapeutic outcome in 2L squamous NSCLC in recent years

1. CancerMPact; 2. Clarivate / Clarivate Survey]

# Data Support Initiation of Pivotal Phase 3 Trial Evaluating Gotistobart<sup>1</sup> in CPI-resistant NSCLC


**PRESERVE-001: Phase 1/2 multicenter, non-randomized, open-label, multiple-dose, FIH trial**  
 He K. et al. ASCO 2023 #9024.

**Anti-tumor activity observed in ICI-resistant NSCLC patients (n=27)**


**ORR: 29.6%** (22.2% confirmed & 7.4% unconfirmed)  
**DCR: 70.4%**

**Manageable adverse events**

**Target lesion best overall response (n=27 evaluable)**  
 Dosing 10 mg/kg x 2, then 6 mg/kg, q3w (2 pts.: 10 mg/kg x 4, q3w)




**Target lesion percentage change over time (n=27 evaluable)**  
 Dosing: 10 mg/kg x 2, then 6 mg/kg, q3w (2 pts.: 10 mg/kg x 4, q3w)



<sup>1</sup>Partnered with OncoC4; \*NCT04140526.

# Pivotal Development of Gotistobart<sup>1</sup> in 2L Squamous Non-Small Cell Lung Cancer




## Extensive-Stage Small Cell Lung Cancer is a High-Incidence Cancer with Poor Long-term Survival Rates

2030 U.S., EU4, U.K. SCLC incidence<sup>1</sup>

~60k

SCLC staging distribution<sup>2</sup>



High unmet need for ES-SCLC patients as long-term survival outcomes remain very poor

|                                    | Limited-Stage SCLC   | Extensive-Stage SCLC                                |
|------------------------------------|--|---|
| <b>mOS</b>                         | cCRT: ~25 – 30 mos (CONVERT) <sup>3</sup><br>Durva consolidation: 56 mos (ADRIATIC) <sup>4</sup> | Atezo + chemo: 12.3 mos (IMPower133) <sup>5,6</sup> |
| <b>24 mos OS</b>                   | cCRT: ~ 50% (CONVERT) <sup>3</sup><br>Durva consolidation: 68% (ADRIATIC) <sup>4</sup>           | Atezo + chemo: ~ 25% (IMPower133) <sup>5,6</sup>    |
| <b>5-year survival<sup>2</sup></b> | 20%  | 3%  |

1. Incidence from: SEER data for diagnosed SCLC incidence in US; Cancer Research UK; Zentrum für Krebsregisterdaten; Santé Publique; AIOM; EPDATA. 2 Statistics from Dayen et al (2019); CancerMPact® Patient Metrics US & EU5, accessed February 2024. \*Due to limited survival data in EU5, U.S. survival data is reported. 3 Wallis, Gerard M. et al. International Journal of Radiation Oncology, Biology, Physics, Volume 119, Issue 5, 1386 – 1390; 4. Cheng et al., N Engl J Med 2024;391:1313-27. 5 L. Horn et al, N. Engl. J. Med., 379 (2018), pp. 2220-2229; 6 Stephen V. Liu et al., JCO 39, 619-630(2021).

## Expeditious Set Up of Global Phase 2 Trial To Confirm Optimal Dose For Pivotal Development

Fully enrolled Global Phase 2 dose-optimization of punitamig<sup>1</sup> + chemotherapy in patients with 1L/2L SCLC

### Key Inclusion Criteria

- Cohort 1: Untreated ES-SCLC or LS-SCLC with TFI ≤ 6 months since last treatment

Punitamig (30 mg/kg Q3W) + chemotherapy

Punitamig (20 mg/kg Q3W) + chemotherapy

Treatment continued until disease progression or intolerable toxicity

- **Three cohorts** based on previous treatment for patients and chemo option used to combine with punitamig
- Each cohort randomized to receive two different dose levels punitamig with chemo

### Key Endpoints



**Primary:** ORR, safety  
**Secondary:** PFS, OS

1. Partnered with Bristol Myers Squibb; 2. L. Horn et al. N. Engl. J. Med., 379 (2018), pp. 2220-2229; NCT06449209.

# Pumitamid<sup>1</sup> Combined with Chemotherapy Indicated Encouraging Efficacy in 1L ES-SCLC in Phase 2 Study

## Efficacy

Encouraging efficacy observed across treatment lines, including >95% disease control rates in 1L-ESCLC


## Safety Profile

Consistent and manageable safety profile across studies with low discontinuation rates and no new safety concerns

## Consistent Clinical Profile

Activity and safety confirmed in China and global datasets, supporting frontline benefit

| Patient Population | 2L SCLC China<br>IO Naive<br>30 mg/kg Q3W | 2L SCLC China<br>IO Treated<br>30 mg/kg Q3W | 1L ES-SCLC China<br>30 mg/kg Q3W | 1L ES-SCLC Global<br>20 mg/kg Q3W | 1L ES-SCLC Global<br>30 mg/kg Q3W |
|--------------------|---|---|----------------------------------|-----------------------------------|-----------------------------------|
| N                  | 22  | 43  | 48                               | 20                                | 18                                |
| cORR (%)           | 50.0                                      | 37.2  | 85.4                             | 85.0                              | 66.7                              |
| DCR (%)            | 81.8                                      | 90.7  | 97.9                             | 100                               | 100                               |
| mPFS (months)      | 5.5                                       | 5.4   | 6.9                              | 6.3                               | 7.0                               |
| mOS (months)       | 14.7                                      | 14.3  | 16.8                             | -                                 | -                                 |
| Congress           | ELCC 2025                                 |   | ELCC 2025                        | WCLC 2025                         |                                   |




### Benchmark data<sup>2</sup> 1L ES-SCLC

| Regimen       | ORR | mPFS   | mOS     | Study                   |
|---------------|-----|--------|---------|-------------------------|
| Atezo + Chemo | 60% | 5.2 mo | 12.3 mo | IMpower133 <sup>3</sup> |
| Durva + Chemo | 68% | 5.1 mo | 12.9 mo | CASPIAN <sup>4</sup>    |


1. Partnered with Bristol Myers Squibb; 2. This benchmarking is not based on head-to-head trials between BioNTech's investigational candidates and other products or product candidates. Furthermore, definitive conclusions cannot be drawn from cross-trial comparisons or anticipated data, as they may be confounded by various factors, and should be interpreted with caution; 3. Horn et al., New England Journal of Medicine, 2018; 4. Paz-Ares et al., The Lancet, 2019.

# Pumitamid<sup>1</sup> Shows Early Signs Of Durable Antitumor Activity in SCLC

**mDoR: 4.9 months (overall)**



**mPFS: 6.8 months (overall)**



Data cut-off: 07 Aug 2025; median follow-up 28.3 weeks (min, max 3.9, 45.6) overall. Median treatment duration: 25.3 weeks (Q1 12.9, Q3 30.6).

# Global Phase 3 Trial to Establish Pumitamid<sup>1</sup> in ES-SCLC

Phase 3 multi-site, double-blinded, randomized trial of pumitamid in combination with chemotherapy compared to atezolizumab with chemotherapy in previously untreated ES-SCLC



## Key Endpoints




**Primary:** OS  
**Secondary:** PFS

## Benchmark Data for 1L ES-SCLC

| Regimen       | ORR | mPFS   | mOS     | Study                   |
|---------------|-----|--------|---------|-------------------------|
| Atezo + chemo | 60% | 5.2 mo | 12.3 mo | IMpower133 <sup>2</sup> |


1. Partnered with Bristol Myers Squibb; 2. L. Horn et al., New England Journal of Medicine, 2018; NCT06712355.

## BioNTech's Currently Ongoing Trials\* in Lung Cancer





Partnered with: 1. Bristol Myers Squibb; 2. OncoC4; 3. DualityBio; 4. MedLink; \*As of Nov 2025; \*\*being conducted in China

## Our Diverse Lung Cancer Pipeline



1. Synergistic potential; Partnered with 2 DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. OncoC4.

## High B7-H3 Protein Expression Observed in Various Solid Tumors



# BNT324/DB-1311<sup>1</sup> Monotherapy Development Focused on Fifteen Phase 2 Dose Optimization and Expansion Cohorts

Phase 1/2 dose escalation, optimization and expansion evaluating BNT324/DB-1311 in patients with advanced/metastatic solid tumors unselected for B7-H3 expression

### Key Inclusion Criteria

- Advanced or metastatic solid tumor that progressed on/after standard systemic treatments
- ≥1 measurable lesion per RECIST v1.1
- ECOG PS 0–1
- Asymptomatic brain metastases are allowed

### Key Exclusion Criteria

- Prior treatment with B7-H3 therapy
- Prior treatment with TOP1 ADC

### Phase 1: Dose escalation (complete)




BNT324/DB-1311 IV Q3W  
Until disease progression/unacceptable toxicity

### Phase 2: Dose optimization/expansion (ongoing), 6 or 9 mg/kg Q3W

#### Dose optimization

|          |       |
|----------|-------|
| Cohort 1 | SCLC  |
| Cohort 2 | NSCLC |

|           |       |
|-----------|-------|
| Cohort 4  | CRPC  |
| Cohort 14 | PROC  |
| Cohort 9  | HNSCC |
| Cohort 13 | HNSCC |



#### Dose expansion

|           |                    |
|-----------|--------------------|
| Cohort 3  | ESCC               |
| Cohort 5  | Melanoma           |
| Cohort 6  | HCC                |
| Cohort 7  | Cervical cancer    |
| Cohort 8  | Other solid tumors |
| Cohort 10 | Rare tumors        |
| Cohort 11 | Post Lu-177 CRPC   |
| Cohort 12 | Taxane-naïve CRPC  |
| Cohort 15 | DDI Cohort         |

### Key Endpoints



**Primary:** DLT/MTD (Phase 1), safety and ORR (Phase 2)  
**Secondary:** DCR, DOR, PFS, OS, and B7-H3 expression


<sup>1</sup>An intermediate dose of 10.5 mg/kg was evaluated after 2 DLTs occurred with 12 mg/kg.  
1. Partnered with DualityBio; Cheng Y et al. ESMO Asia 2024 #570 NCT05914116

Early Signs Of Encouraging Activity With BNT324/DB-1311<sup>1</sup> in NSCLC

Cheng Y et al. ESMO Asia 2024 570.

|                               | Non-squamous NSCLC (n=41) | Squamous NSCLC (n=25)     |
|-------------------------------|---------------------------|---------------------------|
| <b>ORR, n (%)</b><br>[95% CI] | 9 (22.0)<br>[10.6, 37.6]  | 4 (16.0)<br>[4.5, 36.1]   |
| Confirmed ORR, n (%)          | 5 (12.2)                  | 0                         |
| Pending confirmation, n       | 3                         | 4                         |
| <b>DCR, n (%)</b><br>[95% CI] | 33 (80.5)<br>[65.1, 91.2] | 15 (60.0)<br>[38.7, 78.9] |
| <b>3-month PFS rate, %</b>    | 74.5                      | 50.5                      |

EGFRm NSCLC (n=14) ORR: 3 (21.4%)




1. Partnered with DualityBio.

— Encouraging Antitumor Activity BNT324/DB-1311<sup>1</sup> in SCLC

Higher ORR with 9 mg/kg in patients who received prior IO but no prior TOP1i (79% of the patients)

|                                  | SCLC<br>6 mg/kg<br>(n=33) | SCLC<br>9 mg/kg<br>(n=34) |
|----------------------------------|---------------------------|---------------------------|
| <b>ORR, n (%)</b><br>[95% CI]    | 18 (54.5)<br>[36.4, 71.9] | 20 (58.8)<br>[40.7, 75.4] |
| Confirmed ORR, n (%)             | 9 (27.3)                  | 12 (35.3)                 |
| Pending confirmation, n          | 6                         | 4                         |
| <b>DCR, n (%)</b><br>[95% CI]    | 29 (87.9)<br>[71.8, 96.6] | 31 (91.2)<br>[76.3, 98.1] |
| <b>3-month PFS rate, %</b>       | <b>67.4</b>               | <b>79.3</b>               |
| <b>Prior IO + no prior TOP1i</b> | <b>n=15</b>               | <b>n=27</b>               |
| <b>ORR, n (%)</b>                | 7 (46.7)                  | 19 (70.4)                 |
| Confirmed ORR, n (%)             | 3 (20.0)                  | 11 (40.7)                 |
| Pending confirmation             | 3                         | 4                         |



<sup>1</sup>. Partnered with DualityBio; Cheng Y et al. ESMO Asia 2024 570.

# BNT324/DB-1311<sup>1</sup> Safety Profile

➤ More Grade ≥3 TRAEs with 9 mg/kg, but similarly low rate of TRAEs leading to discontinuation


➤ Gastrointestinal and hematological events, primarily Grade 1–2, were the most common TRAEs

➤ Hematological TRAEs occurred more frequently with 9 mg/kg than with 6 mg/kg

## Overall Population

Parsonson A et al. ASCO 2025 5015

| n (%)                  | Overall* (n=465) | 6 mg/kg (n=119) | 9 mg/kg (n=332) |
|------------------------|------------------|-----------------|-----------------|
| Any TRAE               | 429 (92.3)       | 110 (92.4)      | 306 (92.2)      |
| Grade ≥3 TRAE          | 220 (47.3)       | 34 (28.6)       | 178 (53.6)      |
| TRAE leading to:       |                  |                 |                 |
| Dose reduction         | 71 (15.3)        | 6 (5.0)         | 59 (17.8)       |
| Interruption           | 100 (21.5)       | 16 (13.4)       | 81 (24.4)       |
| Discontinuation        | 30 (6.5)         | 6 (5.0)         | 22 (6.6)        |
| TRAE leading to death† | 2 (0.4)          | 0               | 1 (0.3)         |



ILDs/pneumonitis reported in 5 patients receiving 6 mg/kg and 15 receiving 9 mg/kg, all Grade 1–2 except 2 Grade 3 events in two patients receiving 9 mg/kg.

Data cut-off: 04-Mar-2025

1. Partnered with DualityBio. \*Includes 3 mg/kg (n=4), 10.5 mg/kg (n=4), and 12 mg/kg (n=6). The overall population includes patients with other tumors such as SCLC, NSCLC, ESCC, melanoma, HCC, cervical cancer, and HNSCC. †TRAEs leading to death: pneumonitis/respiratory failure in a patient receiving 10.5 mg/kg and encephalopathy in a patient receiving 9 mg/kg. Parsonson A et al. ASCO 2025 5015.

# Evaluating BNT324/DB-1311<sup>1</sup> in Combination with Punitamig<sup>2</sup> in Patients With Advanced Lung Cancer

Two-part study to evaluate efficacy and safety of a combination therapy with BNT324/DB-1311 and punitamig in patients with advanced lung cancer

### Key Inclusion Criteria


- Histologically or cytologically confirmed unresectable adv./met. SCLC or NSCLC
- Measurable disease (RECIST v1.1)
- ECOG PS 0–1
- Eligible regardless of PD-L1 status

### Key Endpoints



- Primary:** Safety (Part 1 and Part 2 Cohorts 1 and 2), ORR (Part 2, all cohorts)
- Secondary:** PFS, OS (Part 2 all cohorts), ORR (Part 1), safety (Part 2, Cohorts 3–7)

### Part 1: Dose escalation (BOIN Design)




### Part 2: Dose optimization and signal-seeking

| Cohort                           | Indication                           | Ratio | RP2D             |
|----------------------------------|--------------------------------------|-------|------------------|
| <b>Dose optimization cohorts</b> |                                      |       |                  |
| DO1                              | 1L nsq NSCLC AGA-                    | R 1:1 | RP2D, RP2D-1     |
| DO2                              | 2L+ SCLC (post-chemo ± IO)           | R 1:1 | RP2D, RP2D-1     |
| <b>Signal-seeking cohorts</b>    |                                      |       |                  |
| 3                                | 2L+ nsq NSCLC AGA- (post-chemo ± IO) |       | RP2D from Part 1 |
| 4                                | 1L sq NSCLC AGA-                     |       |                  |
| 5                                | 2L+ sq NSCLC AGA- (post-chemo ± IO)  |       |                  |
| 6                                | 2L+ nsq NSCLC AGA+ (post-TKI)        |       |                  |
| 7                                | 1L ES-SCLC                           |       |                  |

Partnered with: 1. Duality Bio; 2. Bristol Myers Squibb; NCT06892548

## Our Diverse Lung Cancer Pipeline




1. Synergistic potential; Partnered with 2 DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. OncoC4.

## Evaluating BNT326/YL202<sup>1</sup> in Patients with Advanced NSCLC and BC

| Key Inclusion Criteria  |   | Dose escalation (n=80):<br>A BOIN dose escalation scheme was used for escalation/ de-escalation decisions, followed by cohort backfill at selected doses. |
|---|---|---|
| <b>Inclusion NSCLC</b><br>Locally advanced/ metastatic disease<br>EGFR-activating mutation (exon 19 deletion or L858R)<br>Previous treatment with 3rd generation EGFR TKI, platinum-based CTx, and anti-PD-L1 antibody (US patients)<br>ECOG PS of 0 to 2 | <b>Inclusion BC</b><br>Unresectable, locally advanced or metastatic disease<br>HR+ and HER2- (IHC 0, 1+, 2+/ISH-)<br>Previous treatment with endocrine therapy combined with CDK4/6 inhibitor and 1-2 lines of CTx<br>ECOG PS of 0 to 2 |   |

| Key Endpoints  |  |
|--|--|
|  <b>Primary:</b> Safety and tolerability, MTD<br><b>Secondary:</b> Tumor activity |  |


  

**BNT326/YL202<sup>1</sup>**  
 Target: HER3  
 Payload: topo I inhibitor  
 DAR = 8

<sup>1</sup>. Partnered with MediLink.

# BNT326/YL2021: Encouraging Activity and Near-Complete Disease Control in Patients with Advanced Disease


FIH Phase 1 study: Clinical activity, best percent change from baseline in target lesion size (n=51\*)  
Cheng, Y. et al. ASCO 2024 #3034.



Data Cut-Off: April 16, 2024

1. Partnered with MedLink. \* One patient had non measurable target lesions at PD due to obstructive atelectasis induced by PD in non-target lesions and is therefore not presented in plot. \*\* CR occurred in patient with target lesion as lymph node that shrank to less than 10 mm. \*\*\*26.9% of patients had response confirmed on a subsequent scan; NCT05653752.

# Broad Phase 2 Study to Evaluate BNT326/YL202<sup>1</sup> as a Monotherapy in Various Cancers



1. Partnered with MedLink, NCT06107686

# Evaluating BNT326/YL202<sup>1</sup> in Combination with Punitamig<sup>2</sup> in Patients with Advanced Non-Small Cell Lung Cancer

600+ patients have been treated with BNT326/YL202; being also evaluated in combination with punitamig in NSCLC across histologies, treatment lines

## Key Inclusion Criteria

- Advanced squamous or non-squamous (all cohorts) NSCLC
- Measurable disease defined by RECIST 1.1
- ECOG PS ≤ 1

### Part 1: Dose expansion

2L+, sq or non-sq NSCLC, AGA-neg/pos., any PD-L1

Optional:  
BNT326/YL202 (DL3) + punitamig IV

BNT326/YL202 (DL2) + punitamig IV

BNT326/YL202 (DL1) + punitamig IV

### Part 2a: Dose expansion

Cohort A: 2L+, sq or non-sq NSCLC, AGA-neg/pos., any PD-L1

Arm 1: BNT326/YL202 (DL1) + punitamig.

Arm 2: BNT326/YL202 (DL2) + punitamig

Cohort B: 1L, sq or non-sq NSCLC, AGA-neg, any PD-L1 (n = 40-80)

Arm 1: BNT326/YL202 (DL1) + punitamig

Arm 2: BNT326/YL202 (DL2) + punitamig

### Part 2b: Dose optimization and contribution of components

Cohort C: 2L+, sq or non-sq NSCLC, AGA-negative or EGFR-activating mutation, any PD-L1., any PD-L1

Arm 1: BNT326/YL202 + punitamig, DL1

Arm 2: BNT326/YL202 + punitamig, DL2

Arm 3 – Monotherapy: BNT326/YL202, DL1 or DL2

Cohort D1: 1L, sq or non-sq NSCLC, AGA-negative PD-L1 ≥50%

Arm 1: BNT326/YL202 (DL2) + punitamig

Arm 2 - SOC: pembrolizumab

Arm 3 – Monotherapy: punitamig

Cohort D2: 1L+, sq or non-sq NSCLC, AGA-negative PD-L1 <50%

Arm 1: BNT326/YL202 (DL2) + punitamig

Arm 2 - SOC: pembrolizumab + CTx

## Key Endpoints



Primary:  
Secondary:


Part 1  
Safety

Part 2a  
ORR and safety

Part 2b  
ORR  
PFS, OS

Partnered with: 1. Medilink; 2. Bristol Myers Squibb; (BNT326-02, NCT07111520)

## Our Diverse Lung Cancer Pipeline



1. Synergistic potential; Partnered with 2 DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. OncoC4.

# BNT116<sup>1</sup>-Induced T-Cell Responses Have Been Observed in NSCLC

Vaccine induced CD4+ and CD8+ T-cell responses observed consistently  
 Öven BB, et. al. AACR 2024 CT051


| Patient | Best vaccine response per patient, cell type and target, measured by IFNγ ELISpot post-IVS |      |         |      |         |      |         |      |         |      |       |      | BOR |    |
|---------|--|------|---------|------|---------|------|---------|------|---------|------|-------|------|-----|----|
|         | CLDN6  |      | KK-LC-1 |      | MAGE-A3 |      | MAGE-A4 |      | MAGE-C1 |      | PRAME |      |     |    |
|         | CD4+   | CD8+ | CD4+    | CD8+ | CD4+    | CD8+ | CD4+    | CD8+ | CD4+    | CD8+ | CD4+  | CD8+ |     |    |
| 03-016  |  |      |         |      |         |      |         |      |         |      |       |      |     | SD |
| 03-013  |  |      |         |      |         |      |         |      |         |      |       |      |     | PR |
| 03-018  |  |      |         |      |         |      |         |      |         |      |       |      |     | PR |

■ Induced T cell response (ex vivo ELISpot)    □ No response (ex vivo ELISpot)

Summary of vaccine responses measured post-IVS by IFNγ ELISpot and response per RECISTv1.1


> 90% of NSCLC patients express ≥ 1 TAA  
 > 60% of NSCLC patients express ≥ 2 TAA

De novo and vaccine-expanded T-cell responses were observed across patients  
 Atmaca A, et. al. SITC 2024 P 1486



1. In collaboration with Regeneron; NCT05142189

# Broad Evaluation of FixVac mRNA Immunotherapy in Combination with ADCs and Immunomodulators in Lung Cancer




1. In collaboration with Regeneron; Partnered with: 2. DualityBio; 3. OncoC4; 4. MedLink; 5. Bristol Myers Squibb; NCT05142189.

# BNT116 Has Shown Clinical Activity as Single Agent & in Combination with Chemo or Anti-PD-1 in Advanced NSCLC in Phase 1 Trial<sup>1</sup>

## BNT116 monotherapy plus cemiplimab add-on from cycle 3

Deme et al. SITC 2023

Progression after PD-(L)1 therapy, platinum-based chemotherapy, and one other systemic therapy




Combination therapy with BNT116 plus cemiplimab is active with DCR of 45% in heavily pre-treated lung cancer patients

## BNT116 plus docetaxel

Öven et al. AACR 2024

Progression after PD-(L)1 therapy and platinum-based chemotherapy




Combination therapy with BNT116 plus docetaxel is active with ORR of 30%, DCR of 85% and mPFS of 4.4 months

## BNT116 plus cemiplimab

Atmaca et al. SITC 2024

NSCLC with PD-(L)1 TPS ≥50% that progressed after PD-(L)1 therapy as first-line or adjuvant therapy



Combination therapy with BNT116 plus cemiplimab is active with DCR of 80% and mPFS of 5.5 months


1. In collaboration with Regeneron; NCT05142189.

# BNT116 Has Shown Clinical Activity in Combination with Anti-PD-1 in Advanced NSCLC in Phase 1 Trial<sup>1</sup>

## BNT116 plus cemiplimab in frail patients

Dziadziuszko et al. AACR 2025

Frail patients unfit for 1L chemotherapy with TPS  $\geq 1\%$  advanced NSCLC




Combination therapy with BNT116 plus cemiplimab is active in frail patients with advanced NSCLC with ORR of 45%, DCR of 75% and mPFS of 9.9 months

## BNT116 plus cemiplimab as consolidation treatment

Atmaca et al. WCLC 2025

Advanced NSCLC patients after concurrent chemoradiotherapy



Combination therapy with BNT116 plus cemiplimab indicated encouraging clinical activity with 12-month OS rate of 95%

1. In collaboration with Regeneron; NCT05142189

## Ongoing and Next Steps | Thoracic Cancer

Establishing pumitamidg as a potential  
**frontline treatment for lung cancer**

### ROSETTA Lung-01<sup>1</sup>

Pumitamidg + chemotherapy  
in 1L ES-SCLC

### ROSETTA Lung-02<sup>1</sup>

Pumitamidg + chemotherapy  
in 1L NSCLC

Exploring novel combinations to identify  
**potential future standards-of-care**

### PRESERVE-003<sup>2</sup>

Gotitobart<sup>2</sup>  
in 2L+ sqNSCLC

Part 1 non-pivotal data to be presented at IASLC-NACLC on December 6, 2025

Providing new options for  
**IO experienced sqNSCLC**

### Pumitamidg<sup>1</sup>/ FixVac + ADCs

in metastatic disease

### Pumitamidg<sup>1</sup> + FixVac

in early disease

Novel combination data to be presented in 2026

Partnered with: 1. Bristol Myers Squibb; 2. OncoC4.

92

BIONTECH

# Coffee Break

15 minutes

BIONTECH


---

# Breast Cancer

BIONTECH


---

## Our Diverse Breast Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb.

## BioNTech's Currently Ongoing Trials\* in Breast Cancer




Partnered with: 1. Bristol Myers Squibb; 2. DualityBio; 3. Medilink \*As of November 2025

## TNBC Patients Face Poor Outcomes Due to Limited Therapeutic Options

2030 U.S., EU4, U.K.  
TNBC incidence<sup>1</sup>

~65k

BC staging distribution<sup>2</sup>



Treatment outcomes vary based on PD-L1 levels in 1L TNBC

|   | PD-L1 CPS < 10 (~ 55%) <sup>3,4</sup>             | PD-L1 CPS ≥ 10 (~ 45%) <sup>3,4</sup>                      |
|---|---|--|
| <b>mOS</b>                                  | <b>Chemo: 15.0 mos</b><br>(KN-355) <sup>4</sup>   | <b>Pembro + chemo: 23.0 mos</b><br>(KN-355) <sup>4</sup>   |
| <b>4-year OS</b>                            | <b>Chemo: ~ 15 – 20%</b><br>(KN-355) <sup>4</sup> | <b>Pembro + chemo: ~ 25 – 30%</b><br>(KN-355) <sup>4</sup> |
| <b>5-year survival Stage IV<sup>2</sup></b> | <b>10%</b>  |  |

1. Incidence from SEER (US); Zentrum für Krebsregisterdaten (DE); Globocan (ES); Sante Publique (FR); AIOM (IT); Cancer Research UK 2. CancerMPact® 2024 Treatment Architecture EU5 and US 3. Danziger N, et al, Oncologist, 2023 Apr 6;28(4):319-326. 4. Cortes, J, et al. N. Engl. J. Med. 2022; 387, 217–226.

# Pumitamidg<sup>1</sup> in 1L Triple Negative Breast Cancer

Clinical Benefit Irrespective of PD-L1 Status


Manageable Safety Profile

Novel Combinations Being Evaluated in Parallel

| Patient Population | ITT2 | PD-L1 CPS<1 | PD-L1 1≤CPS<10 | PD-L1 CPS≥10 |
|--------------------|------|-------------|----------------|--------------|
| N                  | 42   | 13          | 16             | 9            |
| cORR (%)           | 73.8 | 76.9        | 56.3           | 100.0        |
| DCR (%)            | 95.2 | 100.0       | 93.8           | 100.0        |
| mPFS (months)      | 13.5 | 18.1        | 14.0           | 10.8         |
| 18-mo OS rate %    | 69.7 | 76.9        | 64.6           | 55.6         |

SABCS 2024

Jiong Wu et al. SABCS 2024 PS3-08



Benchmark data<sup>2</sup> 1L TNBC

| Regimen                   | ORR | mPFS   | mOS     | Study               |
|---------------------------|-----|--------|---------|---------------------|
| Chemo (CPS <10)           | 35% | 5.6 mo | 15.2 mo | KN-355 <sup>3</sup> |
| Pembro + Chemo (CPS ≥ 10) | 53% | 9.7 mo | 23.0 mo |                     |


1. Partnered with Bristol Myers Squibb; 2. This benchmarking is not based on head-to-head trials between BionTech's investigational candidates and other products or product candidates. Furthermore, definitive conclusions cannot be drawn from cross-trial comparisons or anticipated data, as they may be confounded by various factors, and should be interpreted with caution; 3. Cortes, J. et al., New England Journal of Medicine, 2022; \*NCT05918133

# Pumitamig<sup>1</sup> Global Phase 2 in Combination with Chemotherapy for 1L/2L Triple Negative Breast Cancer

Trial enrolling heterogenous population across 1L/2L, IO naïve and experienced

## Key Inclusion Criteria

- Histologically confirmed, Ia/ mTNBC
- 1L or 2L
- If recurrent, stage I-III BC, at least 6 months has elapsed between completion of treatment with curative intent
- ECOG PS 0,1



## Key Endpoints



**Primary:** ORR per RECIST v1.1 and safety according to NCI-CTCAE v5.0

Data expected at SABCS 2025

1. Partnered with Bristol Myers Squibb 2. Cortes, J, et al. N. Engl. J. Med. 2022; BNT327-02; NCT06449222.

# Phase 3 Study of Punitamig<sup>1</sup> in Combination with Chemotherapy in PD-L1 negative TNBC

Phase 3, multi-site, randomized, double-blind trial of punitamig in combination with chemotherapy versus placebo with chemotherapy in previously untreated locally recurrent inoperable or metastatic PD-L1 negative TNBC

### Key Inclusion Criteria

- Locally recurrent inoperable or metastatic TNBC
- Ineligible for PD-(L)1 + chemo per their tumor PD-L1 expression status
- No prior systemic therapy for TNBC in the advanced setting

n=558  
R 1:1

**Punitamig** + chemotherapy of physician's choice

**Placebo** + chemotherapy of physician's choice

### Key Endpoints




**Primary:** PFS (BICR) and OS

### Benchmark Comparator Data for 1L TNBC (CPS <10)

| Regimen      | ORR | mPFS   | mOS     | Benchmark study          |
|--------------|-----|--------|---------|--------------------------|
| Chemotherapy | 35% | 5.6 mo | 15.2 mo | KEYNOTE-355 <sup>2</sup> |


1. Partnered with Bristol Myers Squibb; 2. Cortes, J, et al. N. Engl. J. Med. 2022; NCT07173751.

## Our Diverse Breast Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb.

# Evaluating T-Pam<sup>1</sup> in Patients With Advanced HER2-Expressing Solid Tumors



1. Partnered with DualityBio; NCT05150691.

# Trastuzumab-Pamirtecan<sup>1</sup> Demonstrates Encouraging Antitumor Activity in HER2-Expressing Patients


## Phase 1/2a\*: Clinical Efficacy

Moore K. et al. ASCO 2023 #3023.

Anti-tumor activity in heavily pretreated HER2-expressing patients


|                               | ORR, % | DCR, % |
|-------------------------------|--------|--------|
| All patients (n=52)           | 44.2   | 88.5   |
| HER2+ breast cancer (n=26)    | 50.0   | 96.2   |
| HER2 low breast cancer (n=13) | 38.5   | 84.6   |

HER2 IHC Status: 1+ 2+ 3+



1. Partnered with DualityBio; <sup>†</sup>NCT05150691.


# Phase 3 Trial Design T-Pam<sup>1</sup> in Chemotherapy-Naïve Patients with HR+ HER2-Low Breast Cancer



<sup>1</sup> Partnered with DualityBio; <sup>2</sup> Twelves C. et al. Clinical Breast Cancer. 2021; NCT06018337.


\* Subjects who have received chemotherapy in the neo-adjuv. or adjuv. setting are eligible, as long as they have had a disease-free interval (defined as completion of systemic chemotherapy to diagnosis of adv. or met disease) of >12 months.

# BioNTech's Currently Ongoing Trials\* in Breast Cancer



Partnered with: 1. Bristol Myers Squibb; 2. DualityBio; 3. Medilink \*As of November 2025


## Our Diverse Breast Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink.


# T-Pam: Phase 1/2 Study of T-Pam<sup>1</sup> in Combination with Punitamig<sup>2</sup> in Advanced Breast Cancer

Phase 1/2 trial to evaluate efficacy, safety and pharmacokinetics of T-Pam in combination with punitamig in participants with advanced breast cancer




Partnered with: 1. DualityBio; 2. Bristol Myers Squibb; NCT06827236.

## Our Diverse Breast Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink.

## Phase 2 Trials Evaluating BNT326/YL202<sup>1</sup> as a Monotherapy in Patients with Advanced Breast Cancer



## Ongoing and Next Steps | Breast Cancer

Establishing pumitamidg<sup>1</sup> as a potential **frontline treatment for TNBC**

### ROSETTA Breast-01<sup>1</sup>

Pumitamidg + chemotherapy in 1L TNBC CPS < 10

### China Phase 3

Pumitamidg + chemotherapy in 1L TNBC  
Data expected in 2026

Evaluating novel pumitamidg<sup>1</sup> + ADC combinations to **bring checkpoint inhibition to additional breast cancer subtypes and treatment settings**

### Pumitamidg<sup>1</sup> + ADCs

Novel combination data expected in 2026

Establishing trastuzumab pamirtecana<sup>2</sup> as a **backbone for HER2-expressing breast cancers**

### DYNASTY Breast-02<sup>2</sup>

Trastuzumab pamirtecana in chemo naïve HR+, HER-2 low BC  
Data expected in 2026

Partnered with: 1. Bristol Myers Squibb; 2. DualityBio.

110


BIONTECH

# Gynecologic Cancers

BIONTECH


---

## Our Diverse Gyn Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb.

## BioNTech's Currently Ongoing Trials\* in Gynecologic Cancers



### Registrational trials in gyn cancers

- ▶ Registrational trial with trastuzumab pamirtecán<sup>2</sup> in endometrial cancer

### PoC trials with novel combinations to inform future registrational trials

- ▶ Pumitamig<sup>1</sup> demonstrated encouraging efficacy as monotherapy in gyn cancers
- ▶ Emerging ADC data informed the start of PoC trials for pumitamig<sup>1</sup> + ADC combinations
- ▶ Combination data will inform future registrational trials

Partnered with: 1. Bristol Myers Squibb; 2. DualityBio \*As of November 2025

# T-Pam<sup>1</sup> Clinical Activity Across HER2-Expression Levels in Endometrial Cancer

## Phase 1/2 FIH study: Clinical Efficacy


Moore K. et al. Presented at ESGO 2023. Abstract # 430

- HER2 tumor expression of IHC 1, 2 and 3+: 31%, 41% and 25%, respectively. Clinical response observed across HER2-expression levels, including IHC 1+.
- Patients received median 2 lines of prior treatment. ~60% of patients had prior IO, ~38% prior anti-HER2 antibody.

| Response <sup>a</sup>           | Dose Escalation            |                            | Dose Expansion             | Total (n=17) <sup>b</sup> |
|---------------------------------|----------------------------|----------------------------|----------------------------|---------------------------|
|                                 | 7 mg/kg (n=4) <sup>b</sup> | 8 mg/kg (n=4) <sup>b</sup> | 8 mg/kg (n=9) <sup>b</sup> |                           |
| <b>Unconfirmed ORR, n (%)</b>   | <b>2 (50)</b>              | <b>4 (100)</b>             | <b>4 (44)</b>              | <b>10 (59)</b>            |
| Confirmed ORR, n (%)            | 1 (25)                     | 3 (75)                     | 0                          | 4 (24)                    |
| Pending confirmation ORR, n (%) | 1 (25)                     | 1 (25)                     | 4 (44)                     | 6 (35)                    |
| <b>Unconfirmed DCR, n (%)</b>   | <b>4 (100)</b>             | <b>4 (100)</b>             | <b>8 (89)</b>              | <b>16 (94)</b>            |

<sup>a</sup> By investigator. <sup>b</sup> Response-evaluable subjects, which includes subjects with ≥1 postbaseline overall response.


| Benchmark Data for 2L+ Endometrial Cancer |     |        |         |                          |
|---|-----|--------|---------|--------------------------|
| Regimen                                   | ORR | mPFS   | mOS     | Study                    |
| Single-agent chemo                        | 15% | 3.8 mo | 11.4 mo | KEYNOTE-775 <sup>2</sup> |



Data cut-off: May 8, 2023

1. Partnered with DualityBio; 2. Makker V. et al. NEJM 2022 (ITT population), NCT05150691

# Phase 3 Trial of T-Pam<sup>1</sup> vs Chemotherapy in 2L+ HER2-Expressing Endometrial Cancer



1. Partnered with DualityBio; 2. Makker V, et al. NEJM 2022; NCT06340568

# Evaluating BNT325/DB-1305<sup>1</sup> as a Monotherapy in Platinum Resistant Ovarian Cancer

A multicohort, first-in-human Phase 1/2 trial evaluating BNT325/DB-1305<sup>1</sup> in patients with advanced/metastatic solid tumors unselected for TROP2 expression

### Key Inclusion Criteria

- ≥18 years of age
- ≥1 measurable lesion per RECIST v1.1
- ECOG PS 0–1
- Adequate organ function
- Asymptomatic brain metastases are allowed

### Key Inclusion Criteria

- Ovarian cancer, primary peritoneal cancer, or fallopian tube cancer (high-grade serous histology)
- 1–4 prior lines of systemic therapy
- PROC disease

### Phase 1: Dose escalation

n=18

2 mg/kg up to 5 mg/kg IV, Q3W

Exposure-response (E-R) relationships for efficacy and safety were performed to support dose identification

### Phase 2a: Dose optimization Cohort 3 in 2L–5L PROC

Dose optimization started with 4 mg/kg and was changed per SMC guidance to 3.5 mg/kg after 10 patients had received 4 mg/kg

n=20  
R 1

4 mg/kg Q3W

3 mg/kg Q3W

n=20  
R 2

3.5 mg/kg Q3W

3 mg/kg Q3W

- 58 patients with ovarian cancer received ≥1 dose of DB-1305/BNT325
- 3 mg/kg (n=30) dose identified for further expansion

### Key Endpoints






**Primary:** ORR, safety  
**Secondary:** DCR, DOR, PFS, OS

<sup>1</sup>. Partnered with DualityBio.

# BNT325/DB-1305<sup>1</sup> Shows Durable Antitumor Activity in Previously Treated Ovarian Cancer

Encouraging antitumor activity in previously treated ovarian cancer

Responders\* (n) 8  
mDOR (months) 7.3




Overall mPFS of 7.4 months

Data cut-off: 15-Dec-2024

1. Partnered with DualityBio; \*BOR not included for 3 patients (3 mg/kg). In 2 mg/kg (n=2): 1 PR and 1 PD and in 5 mg/kg (n=2): 1 CR and 1 SD; \*DOR in patients with confirmed response. Rubinstein et al SGO 2025

## BNT324/DB-1311<sup>1</sup> Shows Emerging Activity in Melanoma, HCC, Cervical Cancer and HNSCC

|                           | CC                       | HCC                       | HNSCC                  | Melanoma                 |
|---------------------------|--------------------------|---------------------------|------------------------|--------------------------|
| Treated, n                | 7                        | 12                        | 7                      | 12                       |
| Evaluable for efficacy, n | 4                        | 12                        | 3                      | 11                       |
| ORR, n (%)<br>[95% CI]    | 3 (75.0)<br>[19.4, 99.4] | 3 (25.0)<br>[5.5, 57.2]   | 3 (100)<br>[29.2, 100] | 4 (36.4)<br>[10.9, 69.2] |
| Confirmed ORR, n (%)      | 3 (75.0)                 | 2 (16.7)                  | 2 (66.7)               | 2 (18.2)                 |
| Pending confirmation, n   | 0                        | 1                         | 1                      | 2                        |
| DCR, n (%)<br>[95% CI]    | 4 (100)<br>[39.8, 100]   | 11 (91.7)<br>[61.5, 99.8] | 3 (100)<br>[29.2, 100] | 9 (81.8)<br>[48.2, 97.7] |



1. Partnered with DualityBio; All patients received either 6 mg/kg or 9 mg/kg except for 1 patient with SCCHN who received 10.5 mg/kg. The difference between the number of patients treated and those evaluable for efficacy is due to patients still on treatment without a first post-baseline scan. Cheng Y et al. ESMO Asia 2024 570.

## Pumitamig<sup>1</sup> in Combination with Novel ADCs in High Unmet Need Indications

Ongoing, first-in-human Phase 1/2 trial evaluating BNT325/DB-1305<sup>2</sup> and pumitamig in patients with advanced/metastatic solid tumors: DB-1305-O-1001 study

### Key Inclusion Criteria

- ≥18 years of age
- ≥1 measurable lesion per RECIST v1.1
- ECOG PS 0–1
- Adequate organ function
- Asymptomatic brain metastases allowed

### Part 1: Dose escalation



### Part 2: Dose expansion (n=30)

| Cohort | Indication         |
|--------|--------------------|
| PM1    | 1-2L NSCLC AGA-    |
| PM2    | 2L nsq NSCLC AGA+  |
| PM3    | 1L Cervical cancer |
| PM4    | 2-4L PROC          |
| PM5    | 1L TNBC            |

### Key Endpoints



Primary: DLT, safety, ORR


Partnered with: 1. Bristol Myers Squibb; 2. Duality; NCT05438329

# Preliminary Data Combining Pumitamid<sup>1</sup> with ADC Showed a Manageable Safety Profile with Few Overlapping Toxicities and Signs of Clinical Activity


## Phase 1/2: Pumitamid<sup>1</sup> combined with TROP2 ADC BNT325/DB-1305<sup>2</sup> in 2L–4L PROC

Erika Hamilton et al. AACR 2025 P648

TRAEs occurring in ≥10% of patients receiving BNT325/DB-1305 + pumitamid<sup>1</sup> Q3W (N=67)




Waterfall plot for PROC from dose expansion cohort in 2-4L PROC (N=13)




1. Partnered with: 1. Bristol Myers Squibb; 2. Duality; NCT05438329.

## Our Diverse Gyn Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb.

## Evaluating Punitamig<sup>1</sup> Monotherapy in Patients with Advanced CC and PROC






<sup>1</sup> Partnered with Bristol Myers Squibb; NCT05918445

# Pumitami<sup>1</sup> as Monotherapy Showed Encouraging Efficacy Signals in Patients with PROC

## Phase 1/2 trial: Efficacy signals PROC

Wu, L. et al. ASCO 2024 #5524



PROC: N=34 | 7 PR | 16 SD | ORR of 20.6% and DCR of 67.7%



mPFS was 5.5 months

1. Partnered with Bristol Myers Squibb; NCT05918445


# Pumitamid<sup>1</sup> as Monotherapy Showed Encouraging Efficacy Signals in Patients with CC

## Phase 1/2 trial: Efficacy signals CC

Wu, L. et al. ASCO 2024 #5524



**CC: N=45 | 1 CR | 18 PR | 23 SD | ORR was 42.2% and DCR 93.3% | ORR in patients with PD-L1-positive tumors was 52.4%**



**mPFS was 8.3 months**

Progression Free Survival (months)

1. Partnered with Bristol Myers Squibb; \*NCT05918445

## Ongoing and Next Steps | Gynecological Cancer

### Establishing trastuzumab pamirtecan<sup>2</sup> in **HER2-expressing endometrial cancer**

#### Single arm registrational Phase 2

Trastuzumab pamirtecan<sup>2</sup> in 2L+ HER2-expressing EC

BLA submission planned for 2026

Data to be presented in 2026

#### Confirmatory Phase 3

Trastuzumab pamirtecan<sup>2</sup> in 2L+ HER2-expressing EC

### Evaluating ADC monotherapy and novel pumitamig<sup>1</sup> + ADC combinations in **ovarian and cervical cancers**

#### Pumitamig<sup>1</sup> + ADCs

Novel combination data to be presented in 2026

Partnered with: 1. Bristol Myers Squibb; 2. DualityBio.

125


BIONTECH

# Gastrointestinal Cancers

BIONTECH


---

## Our Diverse GI Cancer Pipeline



1. Synergistic potential; Partnered with 2. DualityBio; 3. MedLink; 4. Bristol Myers Squibb; 5. Genmab; 6. Genentech, a member of the Roche Group.

## BioNTech's Currently Ongoing Trials\* in GI Cancers




Partnered with: 1. Bristol Myers Squibb; 2. Genmab; 3. DualityBio; 4. MediLink.\*As of November 2025

MSS CRC: Significant Global Incidence with Unmet Need for New Therapies

2030 U.S., EU4, U.K. CRC incidence<sup>1</sup> **~470k**

CRC staging distribution<sup>2</sup>



Patients with metastatic MSS-CRC have few treatment options and require new therapeutic strategies

|                  | dMMR / MSI-H  | pMMR / MSS   |
|------------------|---|--|
| <b>mOS</b>       | <b>Pembro: 77.5mo</b><br>(KEYNOTE-177) <sup>3</sup> | <b>Bevacizumab / Cetuximab + chemotherapy: ~30mo<sup>4,5</sup></b>   |
| <b>5-year OS</b> | <b>Pembro: ~55%</b><br>(KEYNOTE-177) <sup>3</sup>   | <b>Bevacizumab / Cetuximab + chemotherapy: ~15-25%<sup>4,5</sup></b> |

1. Globocan – Cancer Tomorrow. 2. CancerMPact® 2024 Treatment Architecture EU5 and US. 3. André, Ann Oncol., 2024. 4. Cremolini, Lancet, 2015. 5. Venook, JAMA, 2017

# Phase 2 Signal Seeking Trial of Punitamig<sup>1</sup> in Combination with Chemotherapy in Colorectal Cancer

Phase 2, multicenter, open label trial to evaluate efficacy and safety of punitamig in combination with chemotherapy in 1L MSS or MSI-L/pMMR metastatic CRC

### Key Inclusion Criteria

- Histologically or cytologically confirmed metastatic colorectal cancer
- No dMMR or MSI-H
- No prior systemic anti-tumor therapy for CRC
- Measurable lesions per RECISTS v1.1
- ECOG PS 0 or 1

n=30  
R 1:1

Punitamig DL1 + CTx regimen 1, IV, Q2W

Punitamig DL2 + CTx regimen 1, IV, Q2W

n=10  
R 1:1

Punitamig DL1 + CTx regimen 2, IV, Q2W

Punitamig DL2 + CTx regimen 2, IV, Q2W

### Key Endpoints




**Primary:** ORR, safety  
**Secondary:** DOR, DCR, PFS, OS

<sup>1</sup>. Partnered with Bristol Myers Squibb; NCT07133750

# Phase 2/3 Study with Punitamig<sup>1</sup> in Combination with Chemotherapy in Patients with CRC

Phase 2/3, randomized study to evaluate safety and efficacy of punitamig in combination with CTx vs. Bevacizumab in combination with CTx in participants with previously untreated, unresectable, or metastatic CRC



1. Partnered with Bristol Myers Squibb; NCT07221357


# Gastric Cancer: Poor Outcomes and Restricted Therapies in Biomarker Negative Patients

2030 U.S., EU4, U.K.  
gastric cancer  
incidence<sup>1</sup>

**~75k**

High unmet need for metastatic gastric cancer patients as long-term survival outcomes are very poor and treatment options remain limited to chemotherapy in biomarker negative patients


Staging distribution<sup>2</sup>



|                                 | HER2 positive (~15-20%)  | HER2 negative (~80-85%)  |
|---------------------------------|--|--|
| <b>PD-L1, CPS ≥ 1 (~80%)</b>    | <b>Pembro<sup>1</sup> + trastuzumb<sup>2</sup> + chemo</b><br>mOS: 20.1 mo<br>2-year OS: 41%<br>(KEYNOTE-811) <sup>3</sup> | <b>Nivolumab + chemotherapy</b><br>mOS: 13.8mo<br>(CheckMate-649) <sup>4</sup> |
| <b>PD-L1, CPS &lt; 1 (~20%)</b> | <b>Trastuzumab<sup>2</sup> + chemo</b><br>mOS: 20.4 mo<br>(KEYNOTE-811) <sup>3</sup>                                       | <b>Chemotherapy</b><br>mOS 12.5mo<br>(CheckMate-649) <sup>4</sup>              |

1.1. Globocan – Cancer Tomorrow; 2. CancerMPact® 2024 Treatment Architecture EU5 and US; 3. Janjigian, NEJM 2024; 4. Janjigian, JCO 2024


# Phase 2/3 Study with Punitamig<sup>1</sup> in Combination with Chemotherapy in 1L Gastric Cancer



<sup>1</sup> Partnered with Bristol Myers Squibb; NCT07221149

# Evaluating BNT314/GEN1059<sup>1</sup> in Combination with Punitamig<sup>2</sup> in Patients with Metastatic Colorectal Cancer

Phase 1/2 trial across 482 patients to evaluate combination BNT314/GEN1059 and punitamig and chemotherapy in patients with advanced metastatic CRC



Partnered with: 1. Genmab; 2. Bristol Myers Squibb; 3. Bennouna J. et al. Lancet Oncol. 2013.


# Pancreatic Cancer Patients Have Poor Long-Term Survival Rates and Limited Treatment Options

2030 U.S., EU4, U.K. pancreatic cancer incidence<sup>1</sup>

**~153k**

High unmet need for pancreatic cancer patients as long-term survival outcomes are very poor and treatment options remain limited to chemotherapy


Staging distribution<sup>2</sup>



|                        | Stage I-III + Stage IV resectable                  | Stage IV metastatic  |
|------------------------|--|--|
| <b>mOS</b>             | <b>Chemo: 53.5 mo</b><br>(PRODIGE-24) <sup>3</sup> | <b>GnP: 8-9 mo</b><br>(MPACT) <sup>4</sup><br><b>FOLFIRINOX: ~11 mo</b><br>(PRODIGE-IV) <sup>5</sup> |
| <b>24 mos OS</b>       | <b>70%</b><br>(PRODIGE-24) <sup>3</sup>            | <b>GnP: 10%</b><br>(MPACT) <sup>4</sup><br><b>FOLFIRINOX: 10%</b><br>(PRODIGE-IV) <sup>5</sup>       |
| <b>5-year survival</b> | <b>43%</b><br>(PRODIGE-24) <sup>3</sup>            | <b>3%</b> <sup>6</sup>   |

1.1. Globocan – Cancer Tomorrow; 2. CancerMPact® 2024 Treatment Architecture EU5 and US; 3. Conroy et al., JAMA Oncol, 2022; 4. Von Hoff et al., N Engl J Med, 2013; 5. Conroy et al., N Engl J Med, 2011; 6. CancerMPact® 2024 Treatment Architecture EU5 and US.

Phase 2 Trial with Pumitamid<sup>1</sup> in Combination  
with Chemotherapy in Patients with PDAC



**Key Endpoints**



**Primary:** ORR, safety

**Benchmark Data for 1L PDAC**

| Benchmark regimen | ORR | mPFS   | mOS     | Benchmark study         |
|-------------------|-----|--------|---------|-------------------------|
| mFOLFIRINOX       | 32% | 6.4 mo | 11.1 mo | MPACT <sup>2</sup>      |
| GnP               | 23% | 5.5 mo | 8.7 mo  | PRODIGE-IV <sup>3</sup> |


1. Partnered with Bristol Myers Squibb; 2. Von Hoff et al., N Engl J Med, 2013; 3. Conroy et al., N Engl J Med, 2011;

## Exploring CA19-9-ADC BNT329 To Build Presence in GI Cancers

### CA19-9 as an ADC target<sup>1-3</sup>

- CA19-9 is highly expressed in PDAC and other GI cancers<sup>4</sup>


| Tumor Site               | CA 19-9+ / total (%) |
|--------------------------|----------------------|
| Pancreas                 | 29/31 (94%)          |
| Bile Duct                | 10/11 (91%)          |
| Transitional (bladder)   | 22/29 (76%)          |
| Distal esophagus/stomach | 21/30 (70%)          |
| Colon                    | 36/51 (71%)          |
| Ovary                    | 22/37 (59%)          |
| Endometrium              | 27/44 (61%)          |



### First-in-Human trial with BNT329

#### Key Inclusion Criteria

- No patient pre-selection based on CA19-9 expression
- All comor tumor indications known to express CA 19-9 (PDAC, Gastric, Endometrial, Colorectal cancer)



Adapted from Loy et al. 1993

1. Passerini R, et al. Am J Clin Pathol 2012;138(2):281-7; 2. Data on file; 3. Lee et al. World J Gastrointest Surg. 2020 Dec 27;12(12): 468-490; 4. Loy et al. Am J Clin Pathol.1993;99:726-728

## Ongoing and Next Steps | Gastrointestinal Cancer

Establishing pumitamidg<sup>1</sup> in  
**gastrointestinal cancers**

### ROSETTA CRC<sup>1</sup>

Pumitamidg<sup>1</sup> + chemotherapy  
in 1L MSS-CRC

### ROSETTA Gastric-204<sup>1</sup>

Pumitamidg<sup>1</sup> + chemotherapy  
in 1L HER2+, PD-L1+ gastric

### Signal-seeking Phase 2

Pumitamidg<sup>1</sup> + chemotherapy  
in 1L PDAC

Evaluating novel pumitamidg<sup>1</sup> +  
immunomodulator and **ADC**  
monotherapy in late-stage disease

Pumitamidg<sup>1</sup> +  
immunomodulator

ADCs

Partnered with: <sup>1</sup> Bristol Myers Squibb.

138


BIONTECH

# Genitourinary Cancers

BIONTECH

---

## Our Diverse GU Cancer Pipeline



1. Synergistic potential; Partnered with: 2. DualityBio; 3. Bristol Myers Squibb; 4. OncoC4.

## BioNTech's Current Development in GU Cancer



Partnered with: 1. Bristol Myers Squibb; 2. DualityBio; 3. OncoC4.

# Evaluating Punitamig<sup>1</sup> Monotherapy in Patients with 2L Clear Cell RCC and 1L Non-Clear Cell RCC

Phase 1/2 multiple cohort monotherapy trial to evaluate safety and efficacy of punitamig in patients with advanced solid tumors, including RCC

## Key Inclusion Criteria

- Locally advanced inoperable or metastatic RCC with or without sarcomatoid component
  - ccRCC: progress on prior 1L VEGF TKI +/- IO
  - nccRCC: no prior systemic therapy
- Malignant tumor confirmed by histology or cytology
- Adequate organ function
- $\geq 1$  measurable lesion not been previously treated (RECIST 1.1)
- ECOG 0-1

n = 53

**Punitamig, iv,  
Q2W or Q3W**

Treatment continued until disease progression or intolerable toxicity

## Key Endpoints




**Primary:** ORR  
**Secondary:** DCR, DOR, PFS, safety

<sup>1</sup>. Partnered with Bristol Myers Squibb; NCT05918445




# Pumitamidg<sup>1</sup> Showed Encouraging Anti-Tumor Activity in Patients with 1L nccRCC Across Subtypes


ORR: 36.4%; DCR: 90.9% (n=22)



mPFS: 15.1 months




DOR of up to 24.9 months



1. Partnered with Bristol Myers Squibb; Xinan Sheng, et. al. ESMO 2024 1692P.

## Our Diverse GU Cancer Pipeline



1. Synergistic potential; Partnered with: 2. DualityBio; 3. Bristol Myers Squibb; 4. OncoC4.

## Broad Exploration of BNT324/DB-1311<sup>1</sup>

Phase 1/2, multicenter trial of BNT324/DB-1311 enrolling 465 patients with advanced/metastatic solid tumors unselected for B7H3 expression, including in Prostate Cancer

### Key Inclusion Criteria

- ≥1 measurable lesion per RECIST v1.1 (bone-only disease allowed)
- ECOG PS 0–1
- Adequate organ function
- Progressive mCRPC (serum testosterone <50 ng/dL and PD as defined by PCWG3 criteria)

### Key Exclusion Criteria

- Prior B7H3 targeted therapy
- Prior TOP1 ADC

| Study part/cohort                          | Additional inclusion criteria  | Dose   |
|--|--|--|
| <b>Phase 1</b><br>Dose escalation/backfill |  | 3 mg/kg up to 12 mg/kg IV Q3W  |
| <b>Phase 2:</b>                            |  |  |
| <b>Cohort 4</b><br>(Dose optimization)     | <ul style="list-style-type: none"> <li>• Prior docetaxel; docetaxel rechallenge allowed</li> <li>• Prior NHT</li> </ul>  | <ul style="list-style-type: none"> <li>• 9 mg/kg IV Q3W (n=20)</li> <li>• 6 mg/kg IV Q3W (n=22)</li> </ul> |
| <b>Cohort 11</b><br>(Post Lu-177)          | <ul style="list-style-type: none"> <li>• 1–2 lines of systemic chemotherapy, including docetaxel</li> <li>• Prior NHT</li> <li>• Prior Lu-177 radioligand therapy</li> </ul> | • 6 mg/kg IV Q3W   |
| <b>Cohort 12</b><br>(Taxane-naïve)         | <ul style="list-style-type: none"> <li>• Taxane-naïve; prior (neo)adjuvant use &gt;12 months earlier allowed</li> <li>• Prior NHT</li> </ul>                                 | • 6 mg/kg IV Q3W   |

### Key Endpoints





**Primary:** ORR, safety  
**Secondary:** DCR, DOR, rPFS

<sup>1</sup>. Partnered with DualityBio; BNT324/DB-1311 treatment continued until disease progression/ unacceptable toxicity (treatment beyond progression was allowed); NCT05914116

— Encouraging Efficacy with BNT324/DB-1311<sup>1</sup> in Late-Line mCRPC

|   | Overall<br>(n=73)           | 6 mg/kg<br>(n=38)           | 9 mg/kg<br>(n=33)           |
|---|-----------------------------|-----------------------------|-----------------------------|
| Response evaluable, n                       | 52                          | 24                          | 28                          |
| <b>ORR, (%)</b><br>[95% CI]                 | <b>42.3</b><br>[28.7, 56.8] | <b>41.7</b><br>[22.1, 63.4] | <b>42.9</b><br>[24.5, 62.8] |
| <b>cORR, (%)</b><br>[95% CI]                | <b>30.8</b><br>[18.7, 45.1] | <b>29.2</b><br>[12.6, 51.1] | <b>32.1</b><br>[15.9, 52.4] |
| Pending confirmation, n                     | 5                           | 3                           | 2                           |
| <b>DCR, (%)</b><br>[95% CI]                 | <b>90.4</b><br>[79.0, 96.8] | <b>91.7</b><br>[73.0, 99.0] | <b>89.3</b><br>[71.8, 97.7] |
| <b>mDOR,<sup>†</sup> months</b><br>[95% CI] | <b>ne</b><br>[4.0, ne]      | <b>ne</b><br>[4.2, ne]      | <b>ne</b><br>[4.0, ne]      |
| Evaluable for rPFS, n                       | 68                          | 33                          | 33                          |
| <b>Median rPFS</b>                          |                             |                             |                             |
| Months [95% CI]                             | <b>ne</b> [5.7, ne]         | <b>ne</b> [5.7, ne]         | <b>ne</b> [5.6, ne]         |
| rPFS events, n (%)                          | 14 (20.6)                   | 8 (24.2)                    | 6 (18.2)                    |
| <b>rPFS rate, %</b>                         |                             |                             |                             |
| 6-month                                     | <b>67.7</b>                 | <b>67.1</b>                 | <b>62.7</b>                 |
| 9-month                                     | 58.0                        | 58.7                        | ne                          |



1. Partnered with DualityBio.

## Ongoing and Next Steps | Genitourinary Cancer

Exploring pumitamidig **in GU cancers**

**China Phase 1/2**

Pumitamidig<sup>1</sup> in 2L+ ccRCC and 1L nccRCC

Evaluating ADCs and novel combinations in GU cancers

**ADCs**

Partnered with: 1. Bristol Myers Squibb.

148

BIONTECH



5

# Innovating Early- Stage Cancer Treatment with mRNA Cancer Immunotherapies

Prof. Özlem Türeci, M.D.  
Chief Medical Officer and Co-founder

BIONTECH

# SARS-CoV-2 mRNA Vaccines Sensitize Tumors to CPI


COVID-19 mRNA vaccines are associated with improved survival in patients with NSCLC or metastatic melanoma receiving immunotherapy in study with over 800 patients

- ~25% of these patients received mRNA COVID-19 vaccines within 100 days of initiating immunotherapy
- Demonstrates potential of mRNA vaccines to stimulate innate immunity

## Article SARS-CoV-2 mRNA vaccines sensitize tumours to immune checkpoint blockade


Adam J. Grippin<sup>1,2,3</sup>, Cristiano Marconi<sup>1,2</sup>, Sage Copling<sup>1,2</sup>, Neo Liu<sup>1,2</sup>, Chen Brasel<sup>1</sup>, Cole Morley<sup>1</sup>, Elliana Hwang<sup>1</sup>, Erik Delpit<sup>1</sup>, Min Wang<sup>1</sup>, Ananya Wai<sup>1</sup>, Seung Beom Jeong<sup>1</sup>, Dhruv Kumar Patel<sup>1</sup>, Frances Weisber<sup>1</sup>, Chao Ma<sup>1</sup>, Eden Gaidarova<sup>1</sup>, Andrew Kim<sup>1</sup>, Cheng Zhao<sup>1</sup>, Anna Delmon<sup>1</sup>, Paul Gaudin<sup>1</sup>, Anshu Gulati<sup>1</sup>, Michael R. Rooney<sup>1</sup>, Benjamin R. Schrank<sup>1</sup>, Yifan Wang<sup>1</sup>, Yifan Wu<sup>1</sup>, Enoch Chang<sup>1</sup>, Ramez Khatib<sup>1</sup>, Kyle Dwyer<sup>1</sup>, Jordan Johnson<sup>1</sup>, Nina Hoffman<sup>1</sup>, Gregory Gaudin<sup>1</sup>, Jacob Weir<sup>1</sup>, Julia Rogstad<sup>1</sup>, Diana Amariy<sup>1</sup>, Neelam Thomas<sup>1</sup>, Anika Dohy<sup>1</sup>, Aox Chen<sup>1</sup>, Nihal Copling<sup>1</sup>, Gabriel Ramirez<sup>1</sup>, John Blum<sup>1</sup>, Alicia Bica-Covian<sup>1</sup>, William Daniels<sup>1</sup>, Michael Long<sup>1</sup>, Jeff Lewis<sup>1</sup>, Xiang Wang<sup>1</sup>, Wee Wee Srisaengdeewong<sup>1</sup>, Ais A. Vajpayee<sup>1</sup>, Andrew Johnson<sup>1</sup>, DICKIE Owen<sup>1</sup>, Jia Lee<sup>1</sup>, Ji Ryan Lee<sup>1</sup>, Ryan Smith<sup>1</sup>, Polina Maitland<sup>1</sup>, Jia Wang<sup>1</sup>, Jiarui Zhang<sup>1</sup>, Don L. Gibbons<sup>1</sup>, Jennifer Hargis<sup>1</sup>, Buffy Y. S. Kuo<sup>1</sup>, John V. Heymach<sup>1</sup>, Hector R. Mendez-Gomez<sup>1</sup>, Wei Jiang<sup>1</sup>, Elise J. Sappour<sup>1,2,3</sup> & Steven H. Lee<sup>1,2,3</sup>

Adam Grippin et al. Nature, 2025; Controlling for 39 covariables with COX proportional hazards regression




# mRNA Immunotherapies for Systemic Delivery and Induction of Potent Polyspecific Immune Responses Against Cancer Antigens


Uridine-based mRNA-lipoplexes (RNA-LPX) administered IV for preferential delivery to the APCs in spleen, lymph nodes and bone marrow



RNA-LPX is optimized for immune response stimulation



RNA-LPX combines targeted antigen delivery with stimulation innate immune signature




Secretory signal peptide for translocation of the nascent polypeptide chain into the endoplasmic reticulum



MITD (MHC class I trafficking domain)

5' cap, UTRs, poly(A) tail engineered for optimized stability and translational performance  
Long single-stranded mRNA format, uridine chemistry and LPX to activate innate immune signature

# Full Exploitation of Cancer Target Space for Induction of Anti-Cancer Immunity




<sup>1</sup> Partnered with Genentech, a member of the Roche Group. <sup>2</sup> Antigens vary across programs; <sup>3</sup> T-cell responses analyzed by ex vivo multimer staining analysis in blood.

## Clinical Trial Execution Across iNeST and FixVac Portfolios

| Individualized immunotherapy: iNeST |  |  |  |  | FixVac  |   |   |
|-------------------------------------|--|--|--|--|---|---|---|
| Autogene cevumeran <sup>1</sup>     |  |  |  |  | BNT111 <sup>2</sup>   | BNT113  | BNT116  |
| Adjuvant                            |  |  | 1L   | R/R  | R/R   | 1L  | Multiple settings   |
| MIUC Phase 2                        | CRC Phase 2  | PDAC Phase 2   | Melanoma Phase 2   | Solid tumors Phase 1   | Melanoma Phase 2  | HPV16+ PDL1+ HNSCC Phase 2/3                      | NSCLC Phase 1 & 2   |
| + Nivolumab                         | Monotherapy  | + Atezolizumab + mFOLFIRINOX   | + Pembrolizumab  | Monotherapy and + Atezolizumab   | + Cemiplimab  | + Pembrolizumab                                   | Mono & combinations including BNT324/DB1311 <sup>2</sup> , BNT326/YL202 <sup>1</sup> and Punitamig <sup>3</sup>   |
| Recruitment ongoing                 | Recruitment ongoing<br>Data presented from epi sub-study at <b>ASCO 2024</b> and from biomarker sub-study at <b>ESMO-GI 2024</b> | Recruitment ongoing<br>Data from Phase 1 trial published in 2023 (Rojas et al., <b>Nature</b> )<br>Follow up data published in February 2025 (Sethna et al., <b>Nature</b> ) | Trial completed (n=125)<br>Primary endpoint (significant PFS improvement) not met. Numerical OS benefit trend observed. Data presented at <b>ESMO 2025</b> | Trial completed (n=272)<br>Data published (Lopez et al., <b>Nature Medicine 2025</b> ) | Trial completed (n=184)<br><b>Positive topline data</b> announced in 2024<br>Data presented at <b>ESMO 2025</b> | Recruitment ongoing<br>Trial updated to Phase 2/3 | Recruitment completed in Phase 2 in 1L NSCLC <sup>2</sup><br>Presented at <b>SITC 2023, AACR 2024</b> and <b>SITC 2024</b> .<br>Data in frail patients presented at <b>AACR 2025</b><br>Data in patients after cCRT presented at <b>WCLC 2025</b> |

<sup>1</sup> Partnered with: 1. Genentech, a member of the Roche Group; 2. In collaboration with Regeneron.

# iNeST<sup>1</sup> Phase 2 in 1L Melanoma – Study Design and Primary endpoint



# Autogene Cevumeran Drives Broad, Durable T-Cell Responses in Majority of Patients



High patient-level immunogenicity: 47/56 (84%) patients mounted  $\geq 1$  ex vivo T-cell response



Breadth, not just presence: Median 3 immunogenic neoantigens per patient (range ~1–13)





Vast majority are *de novo* CD8+ T cell responses and they are of high magnitude



84% "immune responder" rate; 16% had no ELISpot signal at the measured timepoint(s)




Neoantigen specific T-cell clones were detectable up to 1.5 years after last autogene cevumeran dose (median longitudinal follow up: 154 days, range 21 – 559 days).



T-cell clones were detectable up to 1.5 years after last autogene cevumeran dose (median longitudinal follow up: 154 days, range 21 – 559 days).

Sullivan et al. ESMO 2025 #954P

# Autogene Cevumeran<sup>1</sup> Phase 2 in 1L Melanoma



Trend of incremental PFS improvement observed in patients with higher neoantigen response breadth<sup>2</sup>


1. Partnered with Genentech, a member of the Roche Group; 2. Exploratory, non-powered analyses; Group "≤ cut-off" includes ELISpot negative patients

# Autogene Cevumeran<sup>1</sup> Phase 2 Data in 1L Melanoma Yield Insights That Support Current Development Focus



Numerical overall survival trend favoring the combination observed at 12-month and 24-month; no formal testing was performed for this secondary endpoint<sup>2</sup>



A trend of improved OS in patients with immune cell PD-L1 High vs. PD-L1-Low in the autogene cevumeran combination arm vs. the pembrolizumab arm



A trend of improved OS in patients with tumors with low mutation burden treated with autogene cevumeran combination vs. the pembrolizumab arm



Sullivan et al. ESMO 2025 #954P

1. Partnered with Genentech, a member of the Roche Group; 2. The primary analysis occurred after 79 PFS events; the median follow-up time for PFS was 35 months and for OS was 45.1 months in the ITT population.

## Ongoing and Next Steps | mRNA Cancer Immunotherapies

### Evaluating autogene cevumeran<sup>2</sup> in adjuvant-stage disease

#### Phase 2

Autogene cevumeran<sup>2</sup> in adjuvant ctDNA+ stage II (high risk) / stage III MSS-CRC

Update planned for 2026

#### Phase 2

Autogene cevumeran<sup>2</sup> + chemotherapy + atezolizumab in adjuvant PDAC

#### Phase 2

Autogene cevumeran<sup>2</sup> + chemotherapy + nivolumab in adj. MIUC

### Evaluating novel combinations for FixVac

#### Phase 2/3

BNT113 + pembrolizumab in HPV16+, PD-L1+ 1L HNSCC

Data expected in 2026

#### BNT116 + pumitamid<sup>1</sup>

#### BNT116 + gotistobart<sup>3</sup>

#### BNT116 + ADCs

Novel combination in 2026

Partnered with: 1. Bristol Myers Squibb; 2. Genentech, member of the Roche Group; 3. OncoC4.



# 6 BioNTech's Path to Value Creation

Ramón Zapata  
Chief Financial Officer

BIONTECH

## Key 2025 Achievements Position BioNTech for Continued Oncology Innovation and Future Growth

✓ Launched variant-adapted COVID-19 vaccine

✓ Leading COVID-19 market share globally (>50%)

✓ >20 phase 2 and 3 oncology trials ongoing

✓ 30+ novel-novel combination cohorts ongoing across tumors

✓ Completed Biotheus acquisition

✓ Strategic BMS partnership to maximize pumitamid

✓ Increased 2025 revenue guidance to €2.6-2.8 billion<sup>1</sup>

✓ €16.7 billion in cash, cash equivalents and securities<sup>2</sup>

**Maintained Leadership in the COVID-19 Space**

**Advanced Key Oncology Pan-Tumor Programs and Clinical Execution**

**Executed Key Strategic Partnerships and Acquisitions**


**Strengthened Financial Position to Drive Sustained Innovation**

<sup>1</sup> BioNTech increased revenue guidance and now expects its revenues for the full 2025 financial year to be in the range of €2,600 - €2,800 million, from previous range of €1,700 - €2,200 million; please refer to 3Q25 earnings press release and quarterly report on Form 6-K for risks and uncertainties. <sup>2</sup> As of September 30, 2025.

160

**BIONTECH**

Driving Impact with Expansion of Later-Stage Oncology Pivotal Trials, While Maintaining R&D Expenditure



<sup>1</sup> YTD October 31, 2025. Visualization illustrative and not to scale.


— Financial Levers Enable BioNTech's Dynamic R&D Investment

**R&D  
Investment  
Control  
Levers**

- ▶ **Active portfolio management strategy** sets high bar for late-stage investment and balance with high risk/reward programs
- ▶ **Innovative, tailored partnerships** to advance priority programs cost effectively, while strengthening P&L
- ▶ **Early-stage programs empowered** with dedicated budgets and opportunistic biotech in-licensing agreements

Innovative BMS Partnership Structured to Accelerate and Maximize Punitamig,  
While Strengthening BioNTech Short- and Long-Term P&L

**BIONTECH** | **Bristol Myers Squibb®**



- Advancing trials in 10+ indications, including registrational trials, and plans to initiate in 1L microsatellite stable CRC and 1L gastric cancer
- 50/50 partnership and cost sharing structure de-risks R&D activities
- \$3.5 billion up-front and non-contingent payments + \$7.6 billion in milestone payments

**Maximizing potential of next-generation PD-L1xVEGF-A bispecific antibody, punitamig, with global co-development and co-commercialization BMS partnership**

— BioNTech Key Principles on the Path to Value Creation



**Strategic Portfolio Management: Shift to Later-Stage De-risked Programs**

Shifting pipeline focus to later-stage programs with higher POS and de-risk through partnerships



**Optimizing Productivity: Do More with Less Through Operational Excellence**

Focus resources on highest priority programs, optimize cost base to support sustainable development trajectory



**Financial Efficiency: Ensuring Cash Runway and Capital Resilience**

Strong financial position provides strategic optionality



**Speed and Scalability: Commercial Readiness and First-to-Market Approach**

Ensure operational agility and organizational readiness to scale rapidly, build commercial infrastructure, prioritize speed to market

**BioNTech Operating  
from Position of Strength**

**2026**

Key Areas of Focus

**1**

**Combination Therapy Momentum**

Anticipate additional datasets from novel-novel combination trials with pumitamidg

**2**

**Modalities to Disease Areas**

2026 marks BioNTech's movement to a focused disease area specific approach

**3**

**Late-Stage Acceleration**

Expect key late-stage data readouts for initial wave of oncology assets

BIONTECH

## Q&A Panel Discussion



**Prof. Uğur Şahin, M.D.**  
Chief Executive Officer  
Co-founder



**Prof. Özlem Türeci, M.D.**  
Chief Medical Officer  
Co-founder



**Ramón Zapata**  
Chief Financial Officer



**Annemarie Hanekamp**  
Chief Commercial Officer



**Prof. Ilhan Celik, M.D.**  
Vice President, Clinical Development



**Michael Wenger, M.D.**  
Vice President, Clinical Development

Thank you


BIONTECH

# Appendix

BIONTECH

---

# BioNTech's Oncology Pipeline – Phase 2 and Phase 3 Clinical Trials



Partnered with 1. Bristol Myers Squibb; 2. Genentech, member of Roche Group; 3. DualityBio; 4. OncoC4; 5. In collaboration with Regeneron; 6. Trial ongoing in China.

# BioNTech's Oncology Pipeline – Phase 1 and Phase 1/2 Clinical Trials

| Phase 1   | Phase 1/2   |   |
|---|---|---|
| <ul style="list-style-type: none"> <li><b>BNT314/GEN1059<sup>2</sup></b> (EpCAMx4-1BB)<br/>Multiple solid tumors</li> <li><b>BNT317</b><br/>Multiple solid tumors</li> <li><b>BNT326/YL202<sup>5</sup></b> (HER3)<br/>Multiple solid tumors</li> <li><b>BNT211</b> (CLDN6)<br/>Multiple solid tumors</li> <li><b>BNT116</b><br/>Adv. NSCLC</li> </ul> | <ul style="list-style-type: none"> <li><b>Pumitamig<sup>1</sup></b> (PD-L1 x VEGF-A)<br/>1L TNBC<sup>5</sup></li> <li><b>Pumitamig<sup>1</sup></b> (PD-L1 x VEGF-A)<br/>Multiple solid tumors<sup>6</sup></li> <li><b>BNT312/GEN1042<sup>2</sup></b> (CD40x4-1BB)<br/>Multiple solid tumors</li> <li><b>Gotistobart<sup>4</sup></b> (CTLA-4)<br/>mCRPC, + radiotherapy</li> <li><b>Gotistobart<sup>4</sup></b> (CTLA-4)<br/>Multiple solid tumors</li> <li><b>Pumitamig<sup>1</sup> + BNT3213 combination</b><br/>1L HCC<sup>5</sup></li> <li><b>Pumitamig<sup>1</sup> + BNT314/GEN1059<sup>2</sup> combination</b><br/>Advanced CRC <b>NEW</b></li> <li><b>Pumitamig<sup>1</sup> +/- BNT3212 combination</b><br/>Multiple solid tumors <b>NEW</b></li> </ul> | <ul style="list-style-type: none"> <li><b>Pumitamig<sup>1</sup> + Trastuzumab-Pamirtecan<sup>3</sup> combination</b><br/>Adv. or metastatic HR+/- HER2-low, ultra-low or null breast cancer</li> <li><b>Pumitamig<sup>1</sup> + BNT324/DB-1311<sup>3</sup> combination</b><br/>Adv. or metastatic HNSCC, HCC, CC, PROC, NSCLC</li> <li><b>Pumitamig<sup>1</sup> + BNT324/DB-1311<sup>3</sup> combination</b><br/>Adv. or metastatic NSCLC or SCLC</li> <li><b>Pumitamig<sup>1</sup> + BNT325/DB-1305<sup>5</sup> combination</b><br/>Multiple solid tumors, PROC, OC, TNBC, NSCLC</li> <li><b>Pumitamig<sup>1</sup> + BNT326/YL202<sup>5</sup> combination</b><br/>Multiple solid tumors <b>NEW</b></li> <li><b>Pumitamig<sup>1</sup> + BNT326/YL202<sup>5</sup> combination</b><br/>Advanced NSCLC <b>NEW</b></li> <li><b>BNT324/DB-1311<sup>3</sup></b> (B7-H3)<br/>Multiple solid tumors</li> <li><b>BNT325/DB-1305<sup>5</sup></b> (TROP2)<br/>Multiple solid tumors</li> <li><b>BNT329</b><br/>Multiple solid tumors <b>NEW</b></li> </ul> |

■ Next generation IO   
 ■ Targeted therapy   
 ■ mRNA immunotherapy

Partnered with: 1. Bristol Myers Squibb; 2. Genmab; 3. DualityBio; 4. Onco C4; 5. MediLink; 6. Trial ongoing in China.

## Upcoming Data Readouts at Medical Conferences in 2025

|                                   | Indication             | Milestone                             | Congress  |
|-----------------------------------|------------------------|---------------------------------------|-----------|
| <b>Pumitamig<sup>1</sup></b>      | 1L/2L TNBC             | Global Phase 2 dose optimization data | SABCS     |
| <b>Gotistobart<sup>2</sup></b>    | 2L sq NSCLC            | Phase 3 Stage 1 data                  | NACLC     |
| <b>BNT324/DB-1311<sup>3</sup></b> | CC and PROC            | Phase 1/2 data                        | ESMO ASIA |
| <b>BNT326/YL202<sup>4</sup></b>   | HR+HER2-null or low BC | Phase 1/2 data                        | SABCS     |

Partnered with: 1. Bristol Myers Squibb; 2. OncoC4; 3. DualityBio; 4. Medlink.

## Select Data Readouts Set BioNTech Up For Catalyst-Rich Period Ahead

|                                     | Indication             | Milestone                                   | Expected Timing |
|-------------------------------------|------------------------|---|-----------------|
| Pumitamidg <sup>1</sup>             | 1L NSCLC               | Global Phase 2 data                         | 2026            |
|                                     | 1L/2L TNBC             | Global Phase 2 dose optimization data       | 2025            |
|                                     | 1L TNBC                | China Phase 3 data                          | 2026            |
| Gotistobart <sup>2</sup>            | 2L sq NSCLC            | Phase 3 Stage 1 data / Phase 3 Stage 2 data | 2025 / 2026     |
| Trastuzumab-Pamirtican <sup>4</sup> | 2L+ HER2-expressing EC | Phase 2 data                                | 2026            |
|                                     | 2L HER2-Low BC         | Global Phase 3 data                         | 2026            |
| BNT324/DB-1311                      | CC and PROC            | Phase 1/2 data                              | 2025            |
| BNT326/YL202                        | HR+HER2-null or low BC | Phase 1/2 data                              | 2025            |
| Autogene cevumeran <sup>2</sup>     | ctDNA+ adj. CRC        | Phase 2 update                              | 2026            |
| FixVac BNT113                       | HPV16+ H&N             | Phase 3 data                                | 2026            |

Partnered with: 1. Bristol Myers Squibb; 2. OncoC4; 3. Genentech, a member of the Roche Group; 4. DualityBio.

## Abbreviation Directory (1)

|         |   |        |   |         |   |
|---------|---|--------|---|---------|---|
| 4-1BB   | CD137   | CCA    | Cholangiocarcinoma                        | EC      | Endometrial cancer  |
| n L     | nth line                                      | cCRT   | Concurrent chemoradiotherapy              | ECOG    | Eastern Cooperative Oncology Group                        |
| AACR    | American Association for Cancer Research      | CDx    | Cluster of differentiation                | EGFR    | Epidermal growth factor receptor                          |
| (bs)AB  | (bispecific) Antibody                         | CDK4/6 | Cyclin-dependent 4/6                      | ELCC    | European Lung Cancer Congress                             |
| ADA     | Anti-drug antibody                            | ChIP   | Chromatin Immunoprecipitation             | EORTC   | Europ. Organisation. for Research and Treatment of Cancer |
| (bs)ADC | (bispecific) Antibody-drug conjugate          | CI     | Confidence interval                       | EpCAM   | Epithelial cell adhesion molecule                         |
| ADCC    | Antibody-dependent cell-mediated cytotoxicity | CLDN6  | Claudin 6                                 | ESCC    | Esophageal squamous cell carcinoma                        |
| ADCP    | Antibody-dependent cellular phagocytosis      | CPI    | Checkpoint inhibitor                      | ESMO    | European Society for Medical Oncology                     |
| adj     | Adjuvant                                      | CPS    | Combined positive score                   | ESMO GI | European Society for Medical Oncology Gastrointestinal    |
| AE      | Adverse event                                 | CR     | Complete response                         | ES-SCLC | Extensive-stage small cell lung cancer                    |
| AGA     | Actionable oncogenic alteration               | CRC    | Colorectal cancer                         | ESO     | Esophageal  |
| AI      | Artificial intelligence                       | CRPC   | Castration resistant prostate cancer      | ET      | Endocrine therapy   |
| ALK     | Anaplastic large-cell lymphoma kinase         | (c)CRT | (Concurrent) Chemoradiation therapy       | EU      | European Union  |
| AST     | Aspartate aminotransferase                    | ctDNA  | Circulating tumor DNA                     | Fab     | Fragment antigen binding                                  |
| ASCO    | American Society of Clinical Oncology         | CTLA   | Cytotoxic T-lymphocyte-associated protein | Fc(R)   | Fragment crystallizable region                            |
| AU      | Absorbance unit                               | ctl    | Control                                   | FDA     | Food and Drug Administration                              |
| AUC     | Area under curve                              | CTX    | Chemotherapy                              | FIH     | First in human  |
| B7-H3   | B7 Homolog 3                                  | DAR    | Drug-antibody ratio                       | FixVac  | Fixed Antigen Vaccine                                     |
| BC      | Breast cancer                                 | DC     | Dendritic cell                            | FoxP3   | Forkhead-Box-Protein P3                                   |
| BICR    | Blinded independent central review            | DCR    | Disease control rate                      | GBM     | Glioblastoma multiforme                                   |
| BLA     | Biologics License Applications                | DDI    | Drug-drug interaction                     | GC/GEJ  | Gastric/Gastro-esophageal junction cancer                 |
| BMS     | Bristol Myers Squibb                          | DFS    | Disease-free survival                     | GI      | Gastrointestinal  |
| BOIN    | Bayesian optimal interval                     | DL     | Dose level                                | GnP     | Gemcitabine plus nab-paclitaxel                           |
| BOR     | Best overall response                         | DLT    | Dose limiting toxicity                    | GrzmB   | Granzyme B  |
| B-RAF   | Serin/Threonin-Kinase                         | dMMR   | Deficient mismatch repair                 | GTEx    | Genotype-Tissue Expression                                |
| BTC     | Biliary tract cancer                          | DNA    | Desoxyribonucleic acid                    | GU      | Genitourinary   |
| C1D1    | Cycle 1 day 1                                 | DO     | Dose optimization                         | Gyn     | Gynecological   |
| CA 19-9 | Carbohydrate antigen 19-9                     | DoR    | Duration of response                      | h       | Human   |
| CC      | Cervical cancer                               | EAC    | Esophageal adenocarcinoma                 | H&E     | Hematoxylin and Eosin                                     |

## Abbreviation Directory (2)

|        |   |          |  |          |   |
|--------|---|----------|--|----------|---|
| H&N    | Head and neck   | JCO      | Journal of Clinical Oncology                   | NCI PRO- | National Cancer Institute Patient Reported Outcome Common             |
| HCC    | Hepatocellular carcinoma  | KD       | Dissociation constant                          | CTCAE    | Terminology Criteria for Adverse Events                               |
| HER2/3 | Human epidermal growth factor receptor 2/3                        | KK-LC-1  | Kita-Kyushu lung cancer antigen 1              | NCI SEER | National Cancer Institute Surveillance, Epidemiology, and End Results |
| HLA    | Human leukocyte antigen   | LALA     | IgG1 variant L234A/L235A                       | NCT      | National clinical trial   |
| HNSCC  | Head and neck squamous cell carcinoma                             | LDH      | Lactate dehydrogenase                          | NE       | Not evaluable for response  |
| HPV    | Human papilloma virus   | LOD      | Limit of detection                             | NEJM     | The New England Journal of Medicine                                   |
| HR     | Hormone receptor  | LPX      | Lipoplex                                       | NEN      | Neuroendocrine neoplasm   |
| IASLC  | International Association for the Study of Lung Cancer            | LUAD     | Lung adenocarcinoma                            | NGS      | Next generation sequencing  |
| IC     | Immune checkpoint   | LUSC     | Lung squamous carcinoma                        | NHT      | Novel hormonal therapy  |
| IC50   | Half maximal inhibitory concentration                             | MAGE-A3  | Melanoma antigen A3                            | NIH      | National Institutes of Health   |
| ICI    | Immune checkpoint inhibitor                                       | MEKi     | Mitogen-activated protein kinase kinase        | NOD SCID | Non-Obese Diabetic-Severe Combined Immunodeficiency                   |
| IFN    | Interferon  | MHC      | Major histocompatibility complex               | NCT      | National clinical trial   |
| IgG    | Immunoglobulin G  | MITD     | Microtubule interacting and trafficking domain | NE       | Not evaluable for response  |
| IHC    | Immunohistochemistry  | MIUC     | Muscle-invasive urothelial carcinoma           | NEJM     | The New England Journal of Medicine                                   |
| IIT    | Investigator initiated trial                                      | mo       | Months   | NEN      | Neuroendocrine neoplasm   |
| ILD    | Interstitial lung disease   | MOA      | Mechanism of Action                            | NGS      | Next generation sequencing  |
| IL-x   | Interleukin x   | MΦ       | Macrophage                                     | NHT      | Novel hormonal therapy  |
| IMDC   | International Metastatic Renal Cell Carcinoma Database Consortium | mono     | Monotherapy                                    | NIH      | National Institutes of Health   |
| iNeST  | Individualized NeoAntigen-Specific Therapy                        | MPM      | Malignant pleural mesothelioma                 | NOD SCID | Non-Obese Diabetic-Severe Combined Immunodeficiency                   |
| IO     | Immuno-oncology   | MRD      | Minimal residual disease                       | NR       | Not reached   |
| Ipi    | Ipilimumab  | mRNA     | Messenger ribonucleic acid                     | NSCLC    | Non-small cell lung cancer  |
| ISH    | In-situ hybridization   | MSI-H(L) | High(low)-frequency microsatellite instability | OC       | Ovarian cancer  |
| ITT    | Intention to treat  | MSS      | Microsatellite stability                       | OP       | Operation   |
| iv     | Intravenously   | MTD      | Maximum tolerated dose                         | (c)ORR   | (confirmed) Objective response rate                                   |
| IvS    | <i>in vitro</i> stimulation                                       | NA       | Not applicable                                 | OS       | Overall survival  |
| JAMA   | Journal of the American Medical Association                       | NACLC    | Nort America conference on Lung Cancer         | OVA      | Ovalbumin   |
|        |   | NCI      | National Cancer Institute                      | P&L      | Profit and loss statement   |
|        |   |          |  | PARP     | Poly (ADP-ribose) polymerase  |

## Abbreviation Directory (3)

|         |  |          |  |              |   |
|---------|--|----------|--|--------------|---|
| PBMC    | Peripheral blood mononuclear cell            | RP2/3D   | Recommended phase 2/3 dose                       | TLR          | Toll-like receptor                              |
| PBS     | Phosphate buffered saline                    | RPL18    | Ribosomal Protein L18                            | TMB-H (or L) | Tumor mutational burden-high or low             |
| PCWG3   | Prostate Cancer Working Group 3              | R/R      | Relapsed/refractory                              | TME          | Tumor microenvironment                          |
| PD      | Progressive disease                          | RT-qPCR  | Real-time quantitative polymerase chain reaction | TNBC         | Triple-negative breast cancer                   |
| PD      | Pharmacodynamics                             | SABCS    | San Antonio Breast Cancer Symposium              | TNF          | Tumor necrosis factor                           |
| PDAC    | Pancreatic ductal adenocarcinoma             | SCCHN    | Squamous cell carcinoma of head and neck         | TOP1         | Topoisomerase I                                 |
| PD-(L)1 | Programmed cell death protein (ligand) 1     | (ES)SCLC | (Extensive/ stage) small cell lung cancer        | TPS          | Tumor proportion score                          |
| pembro  | Pembrolizumab                                | SD       | Standard deviation                               | TPTE         | Transmembrane phosphatase with tensin homology  |
| PFS     | Progression-free survival                    | SD       | Stable disease                                   | TRAE         | Treatment-related adverse event                 |
| pH      | Potentia hydrogenii                          | SEC      | Selenocysteinyl-HRNA                             | Treg         | Regulatory T cell                               |
| Ph x    | (clinical) Phase x                           | SEC      | United States Securities and Exchange Commission | TRON         | Heimholtz Institute for Translational Oncology  |
| PK      | Pharmacokinetics                             | SEER     | Surveillance, epidemiology, and end results      | TROP2        | Trophoblast cell-surface antigen 2              |
| pMMR    | Proficient mismatch repair                   | SEM      | Standard error of the mean                       | TYR          | Tyrosine  |
| PMX     | Pemetrexed                                   | SITC     | Society of Immunotherapy of Cancer               | UC           | Urothelial cancer                               |
| PoC     | Proof of concept                             | SNP      | Single Nucleotide Polymorphism                   | UK           | United Kingdom                                  |
| POS     | Point of sale                                | SoC      | Standard of care                                 | ULN          | Upper limit of normal                           |
| PR      | Partial response                             | SPR      | Surface Plasmon Resonance                        | U.S.         | United States                                   |
| PR      | Progesterone receptor                        | (N)Sq    | (non-)squamous                                   | UTI          | Urinary tract infection                         |
| PRAME   | Preferentially expressed antigen in melanoma | TAA      | Tumor-associated antigen                         | UTR          | Untranslated region                             |
| PROC    | Platinum-resistant ovarian cancer            | TCGA     | The Cancer Genome Atlas                          | VEGF(R) - A  | Vascular endothelial growth factor (receptor) A |
| PVRIG   | Poliovirus receptor-related immunoglobulin   | TCR      | T-cell receptor                                  | VHH          | Heavy chain variable                            |
| R       | Randomized                                   | TEA      | Tissue engineering acoustophoretic               | WCLC         | World Conference of Lung Cancer                 |
| RAS     | Rat sarcoma                                  | TFI      | Treatment-free interval                          | WHO          | World Health Organization                       |
| RCC     | Renal cell carcinoma                         | TIGIT    | T cell immunoreceptor with Ig and ITIM domains   | WT           | Wild type                                       |
| R&D     | Research and development                     | TIL      | Tumor-infiltrating lymphocytes                   | YTD          | Year to date                                    |
| RECIST  | Response Evaluation Criteria in Solid Tumors | TKI      | Tyrosine kinase inhibitor                        |              |   |
| RLT     | Radioligand therapy                          |          |  |              |   |
| RLU     | Relative light units                         |          |  |              |   |

